

HIV Medication Chart

Nucleoside/Nucleotide Analogue Reverse Transcriptase Inhibitors (NRTI)

Emtriva*
(emtricitabine, FTC)



Epivir*†
(lamivudine, 3TC)



Retrovir*†
(zidovudine, AZT, ZDV)



Videx EC*†
(didanosine, ddl)



Viread*
(tenofovir, TDF)



Zerit*†
(stavudine, d4T)




Ziagen*†
(abacavir, ABC)



Protease Inhibitors (PI)

Aptivus*
(tipranavir, TPV)



Crixivan
(indinavir, IDV)



Invirase
(saquinavir hard gel capsules, SQV)



Kaletra*
(lopinavir/ritonavir, LPV/r)



Lexiva*
(fosamprenavir, FPV)



Norvir*
(ritonavir, RTV)




Prezista*
(darunavir, DRV)



Reyataz
(atazanavir, ATV)



Viracept*
(nelfinavir, NFV)



Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTI)

Edurant
(rilpivirine, RPV)



Intelence
(etravirine, ETV)



Rescriptor
(delavirdine, DLV)



Sustiva (efavirenz, EFV)



Viramune*†
(nevirapine, NVP)



FDA Pregnancy Category D

Entry Inhibitors

Fuzeon
(enfuvirtide, T-20)
Fusion Inhibitor



Selzentry
(maraviroc, MVC)
CCR5 Antagonist



Integrase Inhibitors

Isentress▲
(raltegravir, RAL)



Fixed Dose Combinations

Atripla
(TDF+FTC+EFV)



Combivir†
(AZT+3TC)



FDA Pregnancy Category D

Complera
(TDF+FTC+RPV)



Epzicom
(ABC+3TC)




Stribild
(elvitegravir+cobicistat+FTC+TDF)



Trizivir
(AZT+3TC+ABC)



Truvada
(TDF+FTC)



All pills shown in actual size.

Medication brand names appear in bold. Generic names and commonly used abbreviations appear in parentheses.

* Also available in liquid or powder form.



† Generic formulation available.

▲ Chewable form available.

Medication Schedule

Name _____

Date _____

Name of Medication	Number of pills to take each time	Time of day you are to take this medicine		Food Interactions		Side Effects	TOTAL number of pills each day
							
				With Food <input type="checkbox"/>	Without Food <input type="checkbox"/>		
				With Food <input type="checkbox"/>	Without Food <input type="checkbox"/>		
				With Food <input type="checkbox"/>	Without Food <input type="checkbox"/>		
				With Food <input type="checkbox"/>	Without Food <input type="checkbox"/>		
				With Food <input type="checkbox"/>	Without Food <input type="checkbox"/>		
				With Food <input type="checkbox"/>	Without Food <input type="checkbox"/>		

Helpful Hints:

- Refill prescriptions before you run out. Call for refills when you have at least 3 or 4 days left.
- Use cues as a reminder to take your pills (after a meal or favorite TV show, or before bedtime).
- Use reminder aids such as beepers, alarm clocks, and pillboxes. Ask your pharmacist about these.
- Plan ahead (vacations, travel, count out weekly doses).
- Do not stop taking your medications until you have spoken with your health care provider or pharmacist.
- If you have a severe reaction or in case of emergency, contact your health care provider IMMEDIATELY.

Contact Information: Provider _____

Clinic Phone _____ Pharmacy _____ Phone _____

Colorado AIDS Education & Training Center, University of Colorado
Anschutz Medical Campus, 12631 E. 17th Ave., MS A089, Aurora, CO 80045
Tel: 303-724-0646 Fax 303-724-7297 www.coloradoaetc.org

Developed by Lisa Lawrence, MSW; MeriLou Johnson, MSW, MPA; and Steven Johnson, MD

Discontinued Medications or Formulations

Agenerase
(amprenavir, APV)



Fortovase
(saquinavir soft gel capsule, SQV)



HIVID
(zalcitabine, ddC)



Kaletra
(lopinavir/
ritonavir, LPV/r)



Videx
(didanosine, ddl)

