

	<i>% of Respondents<sup>a</sup></i>						
	<i>Geriatricians</i>			<i>Neurologists</i>			
	<i>Mandatory Reporting (n = 86)</i>	<i>Non-mandatory Reporting (n = 45)</i>	<i>p value</i>	<i>Mandatory Reporting (n = 76)</i>	<i>Non-mandatory Reporting (n = 20)</i>	<i>p value</i>	<i>p value<sup>b</sup></i>
Physicians should assess the driving ability of their older drivers more frequently than their middle-aged drivers	86.9	93.3	>0.05	72.0	75.0	>0.05	>0.05
All persons with mild dementia are unsafe to drive	9.3	11.1	>0.05	12.3	36.8	<0.05	<0.001
For every mile driven, older drivers are at significantly higher risk of crashing than middle-aged drivers	77.6	84.4	>0.05	58.7	55.0	>0.05	>0.05
Overall, patients with dementia who drive with a “co-pilot” are safer than those who do not	24.7	29.5	>0.05	44.7	30.0	>0.05	>0.05
I report patients with active driver’s licences to the provincial Department of Motor Vehicles who have:							
Mild cognitive impairment	42.2	24.4	>0.05	28.0	30.0	>0.05	>0.05
Mild Alzheimer disease	90.7	84.1	>0.05	56.0	40.0	>0.05	<0.001
Mild vascular dementia	90.6	88.9	>0.05	54.7	40.0	>0.05	<0.001
Moderate Alzheimer disease	97.7	97.8	>0.05	85.3	50.0	<0.05	<0.001
Moderate vascular disease	97.7	97.8	>0.05	86.7	50.0	<0.05	<0.001
Severe Alzheimer disease	95.3	93.3	>0.05	85.3	55.0	<0.05	0.001
Severe vascular dementia	95.6	95.6	>0.05	86.7	55.0	<0.05	0.001

<sup>a</sup>Percentage who strongly agreed or agreed for attitudes, and who responded always, often, or sometimes for practices.

<sup>b</sup>Difference between all geriatricians and all neurologists.