



Asthma and Allergy
Foundation of America

STUDENT ASTHMA ACTION CARD



ALA
American Lung Association

Name _____	Grade _____	Age _____	ID Photo
Homeroom Teacher _____	Room _____		
Parent/Guardian Name _____ Address _____	Ph. (x) _____ Ph. (w) _____		
Parent/Guardian Name _____ Address _____	Ph. (x) _____ Ph. (w) _____		
Emergency Phone Contact #1 _____	Name _____	Relationship _____	Phone _____
Emergency Phone Contact #2 _____	Name _____	Relationship _____	Phone _____
Physician Treating Student for Asthma _____	Ph. _____		
Other Physician _____	Ph. _____		

EMERGENCY PLAN

Emergency action is necessary when the student has symptoms such as: _____ or has a peak flow reading of _____.

* Steps to take during an asthma episode:

1. Check peak flow.
 2. Give medications as listed below. Student should respond to treatment in 15-20 minutes.
 3. Contact parent/guardian if _____.

 4. Re-check peak flow.
 5. Seek emergency medical care if the student has any of the following:
 - ✓ Cough constantly
 - ✓ No improvement 15-20 minutes after initial treatment with medication and a relative cannot be reached.
 - ✓ Peak flow of _____
 - ✓ Hard time breathing with
 - Chest and neck pulled in with breathing
 - Stooped body posture
 - Struggling or gasping
 - ✓ Trouble walking or talking
 - ✓ Stops playing and can't start activity again
 - ✓ Lips or fingernails are gray or blue
- } If This Happens, Get Emergency Help Now!

* Emergency Asthma Medications

Name	Amount	When to Use
1 _____		
2 _____		
3 _____		
4 _____		

See reverse for more instructions