



Name: _____ D.O.B.: _____

Allergy to: _____

Weight: _____ lbs. Asthma: ☐ Yes (higher risk for a severe reaction) ☐ No

PLACE
STUDENT'S
PICTURE
HERE

For a suspected or active food allergy reaction:

FOR ANY OF THE FOLLOWING
SEVERE SYMPTOMS

- ☐ If checked, give epinephrine immediately if the allergen was definitely eaten, even if there are no symptoms.


LUNG

Short of breath,
wheezing,
repetitive cough


HEART

Pale, blue, faint,
weak pulse, dizzy


THROAT

Tight, hoarse,
trouble breathing/
swallowing


MOUTH

Significant
swelling of the
tongue and/or lips


SKIN

Many hives over
body, widespread
redness


GUT

Repetitive
vomiting or
severe diarrhea


OTHER

Feeling
something bad is
about to happen,
anxiety, confusion

**OR A
COMBINATION
of mild
or severe
symptoms
from different
body areas.**

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. **Use Epinephrine.**

1. **INJECT EPINEPHRINE IMMEDIATELY.**

2. **Call 911.** Request ambulance with epinephrine.

- Consider giving additional medications (following or with the epinephrine):
 - Antihistamine
 - Inhaler (bronchodilator) if asthma
- Lay the student flat and raise legs. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport student to ER even if symptoms resolve. Student should remain in ER for 4+ hours because symptoms may return.

NOTE: WHEN IN DOUBT, GIVE EPINEPHRINE.

MILD SYMPTOMS

- ☐ If checked, give epinephrine immediately for ANY symptoms if the allergen was likely eaten.


NOSE

Itchy/runny nose, sneezing


MOUTH

Itchy mouth


SKIN

A few hives, mild itch


GUT

Mild nausea/discomfort



1. **GIVE ANTIHISTAMINES, IF ORDERED BY PHYSICIAN**

2. Stay with student; alert emergency contacts.

3. Watch student closely for changes. If symptoms worsen, **GIVE EPINEPHRINE.**

MEDICATIONS/DOSES

Epinephrine Brand: _____

Epinephrine Dose: ☐ 0.15 mg IM ☐ 0.3 mg IM

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if asthmatic): _____

PARENT/GUARDIAN AUTHORIZATION SIGNATURE

DATE

PHYSICIAN/HCP AUTHORIZATION SIGNATURE

DATE