

Unsettling Facts of Life: Bacterial Commensalism, Epithelial Adherence, and Inflammatory Bowel Disease

See article on page 44.

A fascinating and unsettling fact of life, perhaps better known to the GASTROENTEROLOGY readership, is our cohabitation with an astonishing diversity and abundance of colonic bacteria. The human large intestine contains hundreds of culturable bacterial species,^{1–3} and morphologic and molecular analysis suggest an equal number of unculturable species.^{4–6} Although the characterization of these bacteria is incomplete, it is clear that there are a lot of them. For individuals on a Western diet, colonic bacteria exceed 50% of fecal solids.⁷ At 10^{13} to 10^{14} organisms, humans on a cellular basis are perhaps only 10% human.

Gut bacteria directly contribute to both energy salvage and metabolic support of the local mucosa, particularly through bacterial hydrolysis and fermentation of carbohydrate and peptide components of the feces.^{8,9} However, commensal bacteria also pose ongoing risks to the host. Bacterial products, particularly cell wall and outer membrane components stimulatory to the innate immune system, may initiate or aggravate mucosal inflammation in the susceptible host.^{10,11} Inflammation, and deficient or unfavorable metabolic products, also contribute to intestinal disorders ranging from flatulence to cancer. The regulation of microbial make-up and activity is thus a critical aspect of gut homeostasis. A major mode of regulation is interbacterial, through processes such as nutrient competition and quorum sensing.¹² In fact, the manipulation of enteric microflora by diet and microbial competition forms the basis of emerging pre- and probiotic therapies.^{13,14}

More subtly, coevolution has yielded reciprocal host-bacterial regulatory mechanisms at the mucosal surface. For example, high-throughput molecular analysis of intestinal colonization by *Bacteroides thetaiotaomicron* reveals striking changes in bacteria metabolism on one hand, and modulation of epithelial functions on the other, including nutrient absorption, mucosal barrier fortification, and xenobiotic metabolism.^{15,16} One aspect of such reciprocal recognition is glycan-based bacterial-epithelial adherence. However, adherence poses profound hazards to the host—excessive bacterial retention, accumulation of toxic or proinflammatory bacterial products, or mucosal invasion. Accordingly, adherence is attenuated through diverse host processes including mucus produc-

tion, secretory IgA, maintenance of local metabolic barriers, antibacterial molecules, and active cellular processes.^{9,16,17}

In this issue of GASTROENTEROLOGY, Swidsinski et al.¹⁸ report a striking disturbance of adherence attenuation in chronic colitis. Using a clever in situ assay for mucosa-associated bacteria, the investigators find that most patients with persistent colitis (inflammatory bowel disease [IBD] and indeterminate colitis) were distinguished from control groups (self-limited colitis and asymptomatic subjects) by more frequent and excessive levels of such bacteria. The bacteria were diverse phylogenetically, and were not distinguishable from the predominant fecal microbiota in healthy individuals. Enteric bacteria play an essential role in human Crohn's disease and model chronic colitis.¹¹ Mutation of a bacterial sensing gene, *NOD2*, accounts for a major genetic locus of Crohn's disease susceptibility.^{19–21} How do the present findings help us better understand the role of bacteria in chronic colitis?

First, a confounding issue in such studies is the effect of disrupted mucosa on local bacterial behavior. At sites of active inflammation, bacteria may encounter dysfunctional or absent epithelium, with gross changes to barriers attenuating invasion and exposure of stromal sites for bacterial adhesion. However, in a careful morphologic and ultrastructural examination, Swidsinski et al.¹⁸ uncover little evidence for these modes of bacterial invasion. In fact, the most abundant bacteria recovered are from intact rather than inflamed mucosal sites. Hence, the phenotype reflects a disorder that appears to directly enhance bacterial-epithelial adhesion.

What mechanisms might account for the excessive bacterial adhesion? Some studies have reported novel adhesive bacteria species in IBD, but these were limited to strains of *E. coli*.^{22–24} In the present work, a diversity of bacterial species share the trait of epithelial adhesion. This could be explained by a primary mucosal disorder, such as impaired host attenuation processes or aberrant epithelial glycan expression.^{25–27} A role for such mucosal dysfunction deserves exploration from cell biology and genetic perspectives, particularly because the phenotype among individuals is heterogeneous within each patient group. However, recent work by Shoemaker et al.²⁸ points to an alternate explanation. Among intestinal microflora, there is remarkably promiscuous sharing of expressible genes and horizontally transmissible ele-

ments.²⁸ One might speculate that the excessive epithelial adhesion by diverse bacterial species in such patients could reflect the exchange and positive selection for a promiscuous adhesin-bearing episome.

Are these adherent bacteria the inducers of mucosal inflammation, or are they nonpathogenic correlates of colitis? The present study does not address this important question, but it is surprisingly challenging to resolve. In murine models of chronic colitis dependent on ambient enteric bacteria, monoassociated bacterial pathogens are only now being identified. In the few cases where representative bacterial species are established (e.g., *B. vulgatus*, *H. hepaticus*, *E. faecalis*), they are highly restricted to individual models, and their bacterial virulence traits are entirely unknown.^{29–32} The present study's design also does not address the potential role of minor microbial components or the major cohort of nonculturable bacteria as disease factors.

Swidsinski et al.¹⁸ describe an apparently global disturbance of host-bacterial interaction in chronic colitis, involving both inflamed and uninvolved mucosa, and diverse bacterial species. The findings reflect a relationship gone awry, but leave open whether host or bacteria is to blame. Confirmation of this striking observation, and delineation of its mechanism, may provide a new perspective on therapeutic strategies to restore the physiologic host-bacterial interaction.

JONATHAN BRAUN

Department of Pathology and Laboratory Medicine
UCLA School of Medicine
Los Angeles, California

References

- Gorbach SL, Nahas L, Lerner PI, Weinstein L. Studies of intestinal microflora. 1. Effects of diet, age, and periodic sampling on numbers of fecal microorganisms in man. *Gastroenterology* 1967;53:845–855.
- Holdeman LV, Good IJ, Moore WEC. Human fecal flora: variation in bacterial composition within individuals and possible effect of emotional stress. *Appl Environ Microbiol* 1976;31:359–375.
- Finegold SM, Sutter VL. Fecal flora in different populations, with special reference to diet. *Am J Clin Nutr* 1978;1:S116–S122.
- Relman DA. The search for unrecognized pathogens. *Science* 1999;284:1308–1311.
- Wilson KH, Blitchington RB. Human colonic biota studied by ribosomal DNA sequence analysis. *Appl Environ Microbiol* 1996;62:2273–2278.
- Tannock GW, Munro K, Harmsen HJ, Welling GW, Smart J, Gopal PK. Analysis of the fecal microflora of human subjects consuming a probiotic product containing *Lactobacillus rhamnosus* DR20. *Appl Environ Microbiol* 2000;66:2578–2588.
- Stephen AM, Cummings JH. The microbial contribution to human faecal mass. *J Med Microbiol* 1980;13:45–56.
- Savage DC. Microbial ecology of the gastrointestinal tract. *Annu Rev Microbiol* 1977;31:107–133.
- Macfarlane GT, Cummings JH. The colonic flora, fermentation, and large bowel digestive function. In: Phillips SF, Pemberton JH, Shorter RG, ed. *The large intestine: physiology, pathophysiology, and disease*. New York: Raven Press, 1991:51–92.
- Beutler B. Autoimmunity and apoptosis: the Crohn's connection. *Immunity* 2001;15:5–14.
- Sartor RB. Review article: How relevant to human inflammatory bowel disease are current animal models of intestinal inflammation? *Aliment Pharmacol Ther* 1997;11(Suppl)3:89–96.
- de Kievit TR, Iglewski BH. Bacterial quorum sensing in pathogenic relationships. *Infect Immun* 2000;68:4839–4849.
- Gionchetti P, Rizzello F, Venturi A, Brigidi P, Matteuzzi D, Bazzocchi G, Poggioli G, Miglioli M, Campieri M. Oral bacteriotherapy as maintenance treatment in patients with chronic pouchitis: a double-blind, placebo-controlled trial. *Gastroenterology* 2000;119:305–309.
- Shanahan F. Inflammatory bowel disease: immunodiagnostics, immunotherapeutics, and ecotherapeutics. *Gastroenterology* 2001;120:622–635.
- Hooper LV, Wong MH, Thelin A, Hansson L, Falk PG, Gordon JI. Molecular analysis of commensal host-microbial relationships in the intestine. *Science* 2001;291:881–884.
- Hooper LV, Gordon JI. Glycans as legislators of host-microbial interactions: spanning the spectrum from symbiosis to pathogenicity. *Glycobiology* 2001;11:1R–10R.
- Godaly G, Bergsten G, Frendeus B, Hang L, Hedlund M, Karpman D, Samuelsson P, Svensson M, Otto G, Wullt B, Svanborg C. Innate defences and resistance to gram negative mucosal infection. *Adv Exp Med Biol* 2000;485:9–24.
- Swidsinski A, Ladhoff A, Pernthaler A, Swidsinski S, Loening-Baucke V, Ortner M, Weber J, Hoffmann U, Schreiber S, Dietel M, Lochs H. Mucosal flora in inflammatory bowel disease. *Gastroenterology* 2002;122:44–54.
- Hampe J, Cuthbert A, Croucher PJ, Mirza MM, Mascheretti S, Fisher S, Frenzel H, King K, Hasselmeier A, Macpherson AJ, Bridger S, Van Deventer S, Forbes A, Nikolaus S, Lennard-Jones JE, Foelsch UR, Krawczak M, Lewis C, Schreiber S, Mathew CG. Association between insertion mutation in NOD2 gene and Crohn's disease in German and British populations. *Lancet* 2001;357:1925–1928.
- Ogura Y, Bonen DK, Inohara N, Nicolae DL, Chen FF, Ramos R, Britton H, Moran T, Karaliuskas R, Duerr RH, Achkar JP, Brant SR, Bayless TM, Kirschner BS, Hanauer SB, Nunez G, Cho JH. A frameshift mutation in NOD2 associated with susceptibility to Crohn's disease. *Nature* 2001;411:603–606.
- Hugot JP, Chamaillard M, Zouali H, Lesage S, Cezard JP, Belaiche J, Almer S, Tysk C, O'Morain CA, Gassull M, Binder V, Finkel Y, Cortot A, Modigliani R, Laurent-Puig P, Gower-Rousseau C, Macry J, Colombel JF, Sahbatou M, Thomas G. Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* 2001;411:599–603.
- Burke DA, Axon AT. Adhesive *Escherichia coli* in inflammatory bowel disease and infective diarrhoea. *BMJ* 1988;297:102–104.
- Giaffer MH, Holdsworth CD, Duerden BI. Virulence properties of *Escherichia coli* strains isolated from patients with inflammatory bowel disease. *Gut* 1992;33:646–650.
- Darfeuille-Michaud A, Neut C, Barnich N, Lederman E, Di Martino P, Desreumaux P, Gambiez L, Joly B, Cortot A, Colombel JF. Presence of adherent *Escherichia coli* strains in ileal mucosa of patients with Crohn's disease. *Gastroenterology* 1998;115:1405–1413.
- Cortfield AP, Myerscough N, Longman R, Sylvester P, Arul S, Pignatelli M. Mucins and mucosal protection in the gastrointestinal tract: new prospects for mucins in the pathology of gastrointestinal disease. *Gut* 2000;47:589–594.
- Rachmilewitz D, Stampler JS, Karmeli F, Mullins ME, Singel DJ, Loscalzo J, Xavier RJ, Podolsky DK. Peroxynitrite-induced rat colitis—a new model of colonic inflammation. *Gastroenterology* 1993;105:1681–1688.

27. Munkholm P. Crohn's disease - Occurrence, course and prognosis - An epidemiologic cohort-study. *Dan Med Bull* 1997;44:287-302.
 28. Shoemaker NB, Vlamakis H, Hayes K, Salyers AA. Evidence for extensive resistance gene transfer among *Bacteroides* spp. and among *Bacteroides* and other genera in the human colon. *Appl Environ Microbiol* 2001;67:561-568.
 29. Rath HC, Herfarth HH, Ikeda JS, Grenther WB, Hamm TE Jr, Balish E, Taurog JD, Hammer RE, Wilson KH, Sartor RB. Normal luminal bacteria, especially bacteroides species, mediate chronic colitis, gastritis, and arthritis in HLA-B27/human beta2 microglobulin transgenic rats. *J Clin Invest* 1996;98:945-953.
 30. Cahill RJ, Foltz CJ, Fox JG, Dangler CA, Powrie F, Schauer DB. Inflammatory bowel disease: an immunity-mediated condition triggered by bacterial infection with *Helicobacter hepaticus*. *Infect Immun* 1997;65:3126-3131.
 31. Kullberg MC, Ward JM, Gorelick PL, Caspar P, Hieny S, Cheever A, Jankovic D, Sher A. *Helicobacter hepaticus* triggers colitis in specific-pathogen-free interleukin-10 (IL-10)-deficient mice through an IL-12- and gamma interferon-dependent mechanism. *Infect Immun* 1998;66:5157-5166.
 32. Rath HC, Schultz M, Freitag R, Dieleman LA, Li F, Linde HJ, Scholmerich J, Sartor RB. Different subsets of enteric bacteria induce and perpetuate experimental colitis in rats and mice. *Infect Immun* 2001;69:2277-2285.
-
- Address requests for reprints to: Jonathan Braun, M.D., Ph.D.,
Department of Pathology and Lab Medicine, UCLA School of Medicine CHS 13-222, Los Angeles, California 90095-1732. e-mail: jbraun@mednet.ucla.edu; fax: (310) 825-5674.
- © 2002 by the American Gastroenterological Association
0016-5085/02/\$35.00
doi:10.1053/gast.2002.31109