

3 Indices and Standard Form

3.1 Index Notation

Here we revise the use of index notation. You will already be familiar with the notation for squares and cubes

$$\begin{array}{l} a^2 = a \times a \\ a^3 = a \times a \times a \end{array}, \text{ and}$$

this is generalised by defining:

$$a^n = \underbrace{a \times a \times \dots \times a}_{n \text{ of these}}$$



Example 1

Calculate the value of:

- (a) 5^2 (b) 2^5 (c) 3^3 (d) 10^4



Solution

$$\begin{aligned} \text{(a)} \quad 5^2 &= 5 \times 5 \\ &= 25 \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad 2^5 &= 2 \times 2 \times 2 \times 2 \times 2 \\ &= 32 \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad 3^3 &= 3 \times 3 \times 3 \\ &= 27 \end{aligned}$$

$$\begin{aligned} \text{(d)} \quad 10^4 &= 10 \times 10 \times 10 \times 10 \\ &= 10\,000 \end{aligned}$$



Example 2

Copy each of the following statements and fill in the missing number or numbers:

$$\text{(a)} \quad 2^{\square} = 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2$$

$$\text{(b)} \quad 9 = 3^{\square}$$

$$\text{(c)} \quad 1000 = 10^{\square}$$

$$\text{(d)} \quad 5^3 = \square \times \square \times \square$$



Solution

(a) $2^7 = 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2$

(b) $9 = 3 \times 3 = 3^2$

(c) $1000 = 10 \times 10 \times 10 = 10^3$

(d) $5^3 = 5 \times 5 \times 5$



Example 3

(a) Determine 2^5 .

(b) Determine 2^3 .

(c) Determine $2^5 \div 2^3$.

(d) Express your answer to (c) in index notation.



Solution

(a) $2^5 = 32$

(b) $2^3 = 8$

(c) $2^5 \div 2^3 = 32 \div 8$
 $= 4$

(d) $4 = 2^2$



Exercises

1. Calculate:

(a) 2^3

(b) 10^2

(c) 3^2

(d) 10^3

(e) 9^2

(f) 3^3

(g) 2^4

(h) 3^4

(i) 7^2

2. Copy each of the following statements and fill in the missing numbers:

(a) $10 \times 10 \times 10 \times 10 \times 10 = 10^{\square}$

(b) $3 \times 3 \times 3 \times 3 = 3^{\square}$

(c) $7 \times 7 \times 7 \times 7 \times 7 = 7^{\square}$

(d) $8 \times 8 \times 8 \times 8 \times 8 = 8^{\square}$

(e) $5 \times 5 = 5^{\square}$

(f) $19 \times 19 \times 19 \times 19 = 19^{\square}$

(g) $6 \times 6 \times 6 \times 6 \times 6 \times 6 \times 6 = 6^{\square}$

(h) $11 \times 11 \times 11 \times 11 \times 11 \times 11 = 11^{\square}$

3. Copy each of the following statements and fill in the missing numbers:

(a) $8 = 2^{\square}$

(b) $81 = 3^{\square}$

(c) $100 = 10^{\square}$

(d) $81 = 9^{\square}$

(e) $125 = 5^{\square}$

(f) $1\,000\,000 = 10^{\square}$

(g) $216 = 6^{\square}$

(h) $625 = 5^{\square}$

4. Is 10^2 bigger than 2^{10} ?

5. Is 3^4 bigger than 4^3 ?

6. Is 5^2 bigger than 2^5 ?

7. Copy each of the following statements and fill in the missing numbers:

(a) $49 = \square^2$

(b) $64 = \square^3$

(c) $64 = \square^6$

(d) $64 = \square^2$

(e) $100\,000 = \square^5$

(f) $243 = \square^5$

8. Calculate:

(a) $2^2 + 2^3$

(b) $2^2 \times 2^3$

(c) $3^2 + 2^2$

(d) $3^2 \times 2^2$

(e) $2^3 \times 10^3$

(f) $10^3 + 2^5$

9. Calculate:

(a) $(3 + 2)^4$

(b) $(3 - 2)^4$

(c) $(7 - 4)^3$

(d) $(7 + 4)^3$

10. Writing your answers in index form, calculate:

(a) $10^2 \times 10^3$

(b) $2^3 \times 2^7$

(c) $3^4 \div 3^2$

(d) $2^5 \div 2^2$

(e) $10^6 \div 10^2$

(f) $5^4 \div 5^2$

11. (a) Without using a calculator, write down the values of k and m .

$$64 = 8^2 = 4^k = 2^m$$

(b) Complete the following:

$$2^{15} = 32\,768$$

$$2^{14} = \boxed{}$$

(KS3/99Ma/Tier 5-7/P1)

3.2 Laws of Indices

There are three rules that should be used when working with indices:

When m and n are positive integers,

$$1. \quad a^m \times a^n = a^{m+n}$$

$$2. \quad a^m \div a^n = a^{m-n} \quad \text{or} \quad \frac{a^m}{a^n} = a^{m-n} \quad (m \geq n)$$

$$3. \quad (a^m)^n = a^{m \times n}$$

These three results are logical consequences of the definition of a^n , but really need a formal proof. You can 'verify' them with particular examples as below, but this is not a proof:

$$\begin{aligned} 2^7 \times 2^3 &= (2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2) \times (2 \times 2 \times 2) \\ &= 2 \times 2 \\ &= 2^{10} \qquad \qquad \qquad (\text{here } m = 7, n = 3 \text{ and } m + n = 10) \end{aligned}$$

or,

$$\begin{aligned} 2^7 \div 2^3 &= \frac{2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2}{2 \times 2 \times 2} \\ &= 2 \times 2 \times 2 \times 2 \\ &= 2^4 \quad (\text{again } m = 7, n = 3 \text{ and } m - n = 4) \end{aligned}$$

Also, $(2^7)^3 = 2^7 \times 2^7 \times 2^7$

$$= 2^{21} \quad (\text{using rule 1}) \quad (\text{again } m = 7, n = 3 \text{ and } m \times n = 21)$$

The proof of the first rule is given below:



Proof

$$\begin{aligned} a^m \times a^n &= \underbrace{a \times a \times \dots \times a}_m \text{ of these} \times \underbrace{a \times a \times \dots \times a}_n \text{ of these} \\ &= \underbrace{a \times a \times \dots \times a \times a \times a \dots \times a}_{(m+n) \text{ of these}} \\ &= a^{m+n} \end{aligned}$$

The second and third rules can be shown to be true for all positive integers m and n in a similar way.

We can see an important result using rule 2:

$$\frac{x^n}{x^n} = x^{n-n} = x^0$$

but $\frac{x^n}{x^n} = 1$, so

$$\boxed{x^0 = 1}$$

This is true for any non-zero value of x , so, for example, $3^0 = 1$, $27^0 = 1$ and $1001^0 = 1$.



Example 1

Fill in the missing numbers in each of the following expressions:

$$(a) \quad 2^4 \times 2^6 = 2^{\square}$$

$$(b) \quad 3^7 \times 3^9 = 3^{\square}$$

$$(c) \quad 3^6 \div 3^2 = 3^{\square}$$

$$(d) \quad (10^4)^3 = 10^{\square}$$



Solution

$$(a) \quad 2^4 \times 2^6 = 2^{4+6} \\ = 2^{10}$$

$$(b) \quad 3^7 \times 3^9 = 3^{7+9} \\ = 3^{16}$$

$$(c) \quad 3^6 \div 3^2 = 3^{6-2} \\ = 3^4$$

$$(d) \quad (10^4)^3 = 10^{4 \times 3} \\ = 10^{12}$$



Example 2

Simplify each of the following expressions so that it is in the form a^n , where n is a number:

$$(a) \quad a^6 \times a^7$$

$$(b) \quad \frac{a^4 \times a^2}{a^3}$$

$$(c) \quad (a^4)^3$$



Solution

$$(a) \quad a^6 \times a^7 = a^{6+7} \\ = a^{13}$$

$$(b) \quad \frac{a^4 \times a^2}{a^3} = \frac{a^{4+2}}{a^3} \\ = \frac{a^6}{a^3} \\ = a^{6-3} \\ = a^3$$

$$(c) \quad (a^4)^3 = a^{4 \times 3} \\ = a^{12}$$



Exercises

1. Copy each of the following statements and fill in the missing numbers:

(a) $2^3 \times 2^7 = 2^{\square}$

(b) $3^6 \times 3^5 = 3^{\square}$

(c) $3^7 \div 3^4 = 3^{\square}$

(d) $8^3 \times 8^4 = 8^{\square}$

(e) $(3^2)^5 = 3^{\square}$

(f) $(2^3)^6 = 2^{\square}$

(g) $\frac{3^6}{3^2} = 3^{\square}$

(h) $\frac{4^7}{4^2} = 4^{\square}$

2. Copy each of the following statements and fill in the missing numbers:

(a) $a^3 \times a^2 = a^{\square}$

(b) $b^7 \div b^2 = b^{\square}$

(c) $(b^2)^5 = b^{\square}$

(d) $b^6 \times b^4 = b^{\square}$

(e) $(z^3)^9 = z^{\square}$

(f) $\frac{q^{16}}{q^7} = q^{\square}$

3. Explain why $9^4 = 3^8$.

4. Calculate:

(a) $3^0 + 4^0$

(b) $6^0 \times 7^0$

(c) $8^0 - 3^0$

(d) $6^0 + 2^0 - 4^0$

5. Copy each of the following statements and fill in the missing numbers:

(a) $3^6 \times 3^{\square} = 3^{17}$

(b) $4^6 \times 4^{\square} = 4^{11}$

(c) $\frac{a^6}{a^{\square}} = a^4$

(d) $(z^{\square})^6 = z^{18}$

(e) $(a^{19})^{\square} = a^{95}$

(f) $p^{16} \div p^{\square} = p^7$

(g) $(p^{\square})^8 = p^{40}$

(h) $q^{13} \div q^{\square} = q$

6. Calculate:

(a) $\frac{2^3}{2^2} + 3^0$

(b) $\frac{3^4}{3^3} - 3^0$

(c) $\frac{5^4}{5^2} + \frac{6^2}{6}$

(d) $\frac{7^7}{7^5} - \frac{5^9}{5^7}$

(e) $\frac{10^8}{10^5} - \frac{5^6}{5^3}$

(f) $\frac{4^{17}}{4^{14}} - \frac{4^{13}}{4^{11}}$

7. Fill in the missing numbers in each of the following expressions:

(a) $8^2 = 2^{\square}$

(b) $81^3 = 9^{\square} = 3^{\square}$

(c) $25^6 = 5^{\square}$

(d) $4^7 = 2^{\square}$

(e) $125^4 = 5^{\square}$

(f) $1000^6 = 10^{\square}$

(g) $81 = \square^4$

(h) $256 = \square^4 = \square^8$

8. Fill in the missing numbers in each of the following expressions:

(a) $8 \times 4 = 2^{\square} \times 2^{\square}$
 $= 2^{\square}$

(b) $25 \times 625 = 5^{\square} \times 5^{\square}$
 $= 5^{\square}$

(c) $\frac{243}{9} = \frac{3^{\square}}{3^{\square}}$
 $= 3^{\square}$

(d) $\frac{128}{16} = \frac{2^{\square}}{2^{\square}}$
 $= 2^{\square}$

9. Is each of the following statements true or false?

(a) $3^2 \times 2^2 = 6^4$

(b) $5^4 \times 2^3 = 10^7$

(c) $\frac{6^8}{2^8} = 3^8$

(d) $\frac{10^8}{5^6} = 2^2$

10. Copy and complete each expression:

$$(a) \quad (2^6 \times 2^3)^4 = (2^{\square})^4 = 2^{\square} \quad (b) \quad \left(\frac{3^6}{3^2}\right)^5 = (3^{\square})^5 = 3^{\square}$$

$$(c) \quad \left(\frac{2^3 \times 2^4}{2^7}\right)^4 = (2^{\square})^4 = 2^{\square} \quad (d) \quad \left(\frac{3^2 \times 9}{3^3}\right)^4 = (3^{\square})^4 = 3^{\square}$$

$$(e) \quad \left(\frac{6^2 \times 6^8}{6^3}\right)^4 = (6^{\square})^4 = 6^{\square} \quad (f) \quad \left(\frac{7^8}{7^2 \times 7^3}\right)^5 = (7^{\square})^5 = 7^{\square}$$

3.3 Negative Indices

Using negative indices produces fractions. In this section we practice working with negative indices. From our work in the last section, we see that

$$a^2 \div a^3 = a^{2-3} = a^{-1}$$

but we know that

$$a^2 \div a^3 = \frac{a \times a}{a \times a \times a} = \frac{1}{a}, \text{ a fraction.}$$

So clearly,

$$a^{-1} = \frac{1}{a}$$

In same way,

$$\begin{aligned} a^{-2} &= \frac{1}{a^2} \\ &= \frac{1}{a \times a} \\ \\ a^{-3} &= \frac{1}{a^3} \\ &= \frac{1}{a \times a \times a} \end{aligned}$$

and, in general,

$$a^{-n} = \frac{1}{a^n}$$

for positive integer values of n . The three rules at the start of section 3.2 can now be used for any integers m and n , not just for positive values.



Example 1

Calculate, leaving your answers as fractions:

(a) 3^{-2}

(b) $2^{-1} - 4^{-1}$

(c) 5^{-3}



Solution

$$\begin{aligned} \text{(a)} \quad 3^{-2} &= \frac{1}{3^2} \\ &= \frac{1}{9} \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad 2^{-1} - 4^{-1} &= \frac{1}{2} - \frac{1}{4} \\ &= \frac{1}{4} \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad 5^{-3} &= \frac{1}{5^3} \\ &= \frac{1}{125} \end{aligned}$$



Example 2

Simplify:

(a) $\frac{6^7}{6^9}$

(b) $6^4 \times 6^{-3}$

(c) $(10^2)^{-3}$



Solution

$$\begin{aligned} \text{(a)} \quad \frac{6^7}{6^9} &= 6^{7-9} \\ &= 6^{-2} = \frac{1}{6^2} = \frac{1}{36} \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad 6^4 \times 6^{-3} &= 6^{4+(-3)} \\ &= 6^{4-3} = 6^1 = 6 \end{aligned}$$

$$\begin{aligned}
 \text{(c)} \quad (10^2)^{-3} &= 10^{-6} \\
 &= \frac{1}{10^6} \\
 &= \frac{1}{1000000}
 \end{aligned}$$



Exercises

1. Write the following numbers as fractions *without using any indices*:

- | | | |
|--------------|--------------|---------------|
| (a) 4^{-1} | (b) 2^{-3} | (c) 10^{-3} |
| (d) 7^{-2} | (e) 4^{-3} | (f) 6^{-2} |

2. Copy the following expressions and fill in the missing numbers:

- | | |
|--|---|
| (a) $\frac{1}{49} = \frac{1}{7^{\square}} = 7^{\square}$ | (b) $\frac{1}{100} = \frac{1}{10^{\square}} = 10^{\square}$ |
| (c) $\frac{1}{81} = \frac{1}{9^{\square}} = 9^{\square}$ | (d) $\frac{1}{16} = \frac{1}{2^{\square}} = 2^{\square}$ |
| (e) $\frac{1}{10000000} = \frac{1}{10^{\square}} = 10^{\square}$ | (f) $\frac{1}{1024} = \frac{1}{2^{\square}} = 2^{\square}$ |

3. Calculate:

- | | |
|------------------------|-------------------------|
| (a) $4^{-1} + 3^{-1}$ | (b) $6^{-1} + 2^{-1}$ |
| (c) $5^{-1} - 10^{-1}$ | (d) $10^{-2} - 10^{-3}$ |
| (e) $4^{-1} - 10^{-1}$ | (f) $6^{-1} + 7^{-1}$ |

4. Simplify the following expressions giving your answers in the form of a number to a power:

- | | |
|--------------------------|--------------------------|
| (a) $4^7 \times 4^{-6}$ | (b) $5^7 \times 5^{-3}$ |
| (c) $\frac{7^4}{7^{-6}}$ | (d) $(3^2)^{-4}$ |
| (e) $(6^{-2})^{-3}$ | (f) $8^4 \times 8^{-9}$ |
| (g) $\frac{7^2}{7^{-2}}$ | (h) $\frac{8^9}{8^{-9}}$ |

5. Copy each of the following expressions and fill in the missing numbers;

(a) $\frac{1}{9} = 3^{\square}$

(b) $\frac{1}{100} = 10^{\square}$

(c) $\frac{1}{125} = 5^{\square}$

(d) $\frac{5}{5^4} = 5^{\square}$

(e) $\frac{6^2}{6^3} = 6^{\square}$

(f) $\frac{2^2}{2^{10}} = 2^{\square}$

6. Simplify the following expressions:

(a) $\frac{x^8}{x^3}$

(b) $\frac{x^7}{x^9}$

(c) $\frac{x^4}{x^8}$

(d) $(x^6)^{-4}$

(e) $\left(\frac{1}{x^2}\right)^4$

(f) $(x^{-8})^3$

7. Copy and complete the following statements:

(a) $0.1 = 10^{\square}$

(b) $0.25 = 2^{\square}$

(c) $0.0001 = 10^{\square}$

(d) $0.2 = 5^{\square}$

(e) $0.001 = 10^{\square}$

(f) $0.02 = 50^{\square}$

8. Copy the following expressions and fill in the missing numbers:

(a) $\frac{x^4}{x^{\square}} = x^2$

(b) $x^6 \times x^{\square} = x^2$

(c) $x^9 \times x^{\square} = x^2$

(d) $\frac{x^7}{x^{\square}} = x^{-2}$

(e) $\frac{x^3}{x^{\square}} = x^4$

(f) $(x^3)^{\square} = x^{-6}$

9. Copy the following expressions and fill in the missing numbers:

(a) $\frac{1}{8} = 2^{\square}$

(b) $\frac{1}{25} = 5^{\square}$

(c) $\frac{1}{81} = 9^{\square}$

(d) $\frac{1}{10000} = 10^{\square}$

10. If $a = b^3$ and $b = \frac{1}{c^2}$, express a as a power of c , without having any fractions in your final answer.

3.4 Standard Form

Standard form is a convenient way of writing very large or very small numbers. It is used on a scientific calculator when a number is too large or too small to be displayed on the screen.

Before using standard form, we revise multiplying and dividing by powers of 10.



Example 1

Calculate:

(a) 3×10^4

(b) 3.27×10^3

(c) $3 \div 10^2$

(d) $4.32 \div 10^4$



Solution

(a) $3 \times 10^4 = 3 \times 10000$
 $= 30\,000$

(b) $3.27 \times 10^3 = 3.27 \times 1000$
 $= 3270$

(c) $3 \div 10^2 = \frac{3}{100}$
 $= 0.03$

$$\begin{aligned}
 \text{(d)} \quad 4.32 \div 10^4 &= \frac{4.32}{10000} \\
 &= \frac{432}{1000000} \\
 &= 0.000432
 \end{aligned}$$

These examples lead to the approach used for standard form, which is a reversal of the approach used in Example 1.

In *standard form*, numbers are written as

$$a \times 10^n$$

where $1 \leq a < 10$ and n is an integer.



Example 2

Write the following numbers in standard form:

- | | |
|-------------|---------------|
| (a) 5720 | (b) 7.4 |
| (c) 473 000 | (d) 6 000 000 |
| (e) 0.09 | (f) 0.000621 |



Solution

- (a) $5720 = 5.72 \times 1000$
 $= 5.72 \times 10^3$
- (b) $7.4 = 7.4 \times 1$
 $= 7.4 \times 10^0$
- (c) $473\,000 = 4.73 \times 100\,000$
 $= 4.73 \times 10^5$
- (d) $6\,000\,000 = 6 \times 1\,000\,000$
 $= 6 \times 10^6$
- (e) $0.09 = \frac{9}{100}$
 $= 9 \div 10^2$
 $= 9 \times 10^{-2}$

$$\begin{aligned}
 \text{(f)} \quad 0.000621 &= \frac{6.21}{10000} \\
 &= \frac{6.21}{10^4} \\
 &= 6.21 \times 10^{-4}
 \end{aligned}$$



Example 3

Calculate:

$$\text{(a)} \quad (3 \times 10^6) \times (4 \times 10^3)$$

$$\text{(b)} \quad (6 \times 10^7) \div (5 \times 10^{-2})$$

$$\text{(c)} \quad (3 \times 10^4) + (2 \times 10^5)$$



Solution

$$\begin{aligned}
 \text{(a)} \quad (3 \times 10^6) \times (4 \times 10^3) &= (3 \times 4) \times (10^6 \times 10^3) \\
 &= 12 \times 10^9 \\
 &= 1.2 \times 10^1 \times 10^9 \\
 &= 1.2 \times 10^{10}
 \end{aligned}$$

$$\begin{aligned}
 \text{(b)} \quad (6 \times 10^7) \div (5 \times 10^{-2}) &= (6 \div 5) \times (10^7 \div 10^{-2}) \\
 &= 1.2 \times 10^9
 \end{aligned}$$

$$\begin{aligned}
 \text{(c)} \quad (3 \times 10^4) + (2 \times 10^5) &= 30000 + 200000 \\
 &= 230000 \\
 &= 2.3 \times 10^5
 \end{aligned}$$



Note on Using Calculators

Your calculator will have a key $\boxed{\text{EE}}$ or $\boxed{\text{EXP}}$ for entering numbers in standard form.

For example, for 3.2×10^7 , press

$$\boxed{3} \quad \boxed{\cdot} \quad \boxed{2} \quad \boxed{\text{EXP}} \quad \boxed{7}$$

which will appear on your display like this:

$$3.2 \times 10^7$$

Some calculators also display the ' $\times 10$ ' part of the number, but not all do. You need to find out what your calculator displays. Remember, you must always write the ' $\times 10$ ' part when you are asked to give an answer in standard form.



Exercises

1. Calculate:

- | | | |
|------------------------|--------------------------|--------------------------|
| (a) 6.21×1000 | (b) 8×10^3 | (c) 4.2×10^2 |
| (d) $3 \div 1000$ | (e) $6 \div 10^2$ | (f) $3.2 \div 10^3$ |
| (g) 6×10^{-3} | (h) 9.2×10^{-1} | (i) 3.6×10^{-2} |

2. Write each of the following numbers in standard form:

- | | |
|--------------------|--------------------|
| (a) 200 | (b) 8000 |
| (c) 9 000 000 | (d) 62 000 |
| (e) 840 000 | (f) 12 000 000 000 |
| (g) 61 800 000 000 | (h) 3 240 000 |

3. Convert each of the following numbers from standard form to the normal decimal notation:

- | | | |
|------------------------|------------------------|------------------------|
| (a) 3×10^4 | (b) 3.6×10^4 | (c) 8.2×10^3 |
| (d) 3.1×10^2 | (e) 1.6×10^4 | (f) 1.72×10^5 |
| (g) 6.83×10^4 | (h) 1.25×10^6 | (i) 9.17×10^3 |

4. Write each of the following numbers in standard form:

- | | |
|---------------|----------------------|
| (a) 0.0004 | (b) 0.008 |
| (c) 0.142 | (d) 0.0032 |
| (e) 0.00199 | (f) 0.000000062 |
| (g) 0.0000097 | (h) 0.00000000000021 |

5. Convert the following numbers from standard form to the normal decimal format:

(a) 6×10^{-2}	(b) 7×10^{-1}	(c) 1.8×10^{-3}
(d) 4×10^{-3}	(e) 6.2×10^{-3}	(f) 9.81×10^{-4}
(g) 6.67×10^{-1}	(h) 3.86×10^{-5}	(i) 9.27×10^{-7}

6. Without using a calculator, determine:

(a) $(4 \times 10^4) \times (2 \times 10^5)$	(b) $(2 \times 10^6) \times (3 \times 10^5)$
(c) $(6 \times 10^4) \times (8 \times 10^{-9})$	(d) $(3 \times 10^{-8}) \times (7 \times 10^{-4})$
(e) $(6.1 \times 10^6) \times (2 \times 10^{-5})$	(f) $(3.2 \times 10^{-5}) \times (4 \times 10^{-9})$

7. Without using a calculator, determine:

(a) $(9 \times 10^7) \div (3 \times 10^4)$	(b) $(8 \times 10^5) \div (2 \times 10^{-2})$
(c) $(6 \times 10^{-2}) \div (2 \times 10^{-3})$	(d) $(6 \times 10^4) \div (3 \times 10^{-6})$
(e) $(4.8 \times 10^{12}) \div (1.2 \times 10^3)$	(f) $(3.6 \times 10^8) \div (9 \times 10^3)$

8. Without a calculator, determine the following, giving your answers in both normal and standard form::

(a) $(6 \times 10^5) + (3 \times 10^6)$	(b) $(6 \times 10^2) + (9 \times 10^3)$
(c) $6 \times 10^5 - 1 \times 10^4$	(d) $8 \times 10^{-2} + 9 \times 10^{-3}$
(e) $6 \times 10^{-4} + 8 \times 10^{-3}$	(f) $6 \times 10^{-4} - 3 \times 10^{-5}$

9. Use a calculator to determine:

(a) $(3.4 \times 10^6) \times (2.1 \times 10^4)$	(b) $(6 \times 10^{21}) \times (8.2 \times 10^{-11})$
(c) $(3.6 \times 10^5) \times (4.5 \times 10^7)$	(d) $(8.2 \times 10^{11}) \div (4 \times 10^{-8})$
(e) $(1.92 \times 10^6) \times (3.2 \times 10^{-11})$	(f) $(6.2 \times 10^{14})^3$

10. The radius of the earth is 6.4×10^6 m. Giving your answers in standard form, correct to 3 significant figures, calculate the circumference of the earth in:

(a) m (b) cm (c) mm (d) km

11. Sir Isaac Newton (1642-1727) was a mathematician, physicist and astronomer.

In his work on the gravitational force between two bodies he found that he needed to multiply their masses together.

- (a) Work out the value of the mass of the Earth multiplied by the mass of the Moon. Give your answer in standard form.

$$\text{Mass of Earth} = 5.98 \times 10^{24} \text{ kg}$$

$$\text{Mass of Moon} = 7.35 \times 10^{22} \text{ kg}$$

Newton also found that he needed to work out the square of the distance between the two bodies.

- (b) Work out the square of the distance between the Earth and the Moon. Give your answer in standard form.

$$\text{Distance between Earth and Moon} = 3.89 \times 10^5 \text{ km}$$

Newton's formula to calculate the gravitational force (F) between two

bodies is $F = \frac{Gm_1m_2}{R^2}$ where

G is the gravitational constant, m_1 and m_2 are the masses of the two bodies, and R is the distance between them.

- (c) Work out the gravitational force (F) between the Sun and the Earth using the formula $F = \frac{Gm_1m_2}{R^2}$ with information in the box below.

Give your answer in standard form.

$$m_1m_2 = 1.19 \times 10^{55} \text{ kg}^2$$

$$R^2 = 2.25 \times 10^{16} \text{ km}^2$$

$$G = 6.67 \times 10^{-20}$$

(KS3/95/Ma/Levels 6-8/P1)



12. (a) Which of these statements is true?
- (i) 4×10^3 is a larger number than 4^3 .
 - (ii) 4×10^3 is the same size as 4^3 .
 - (iii) 4×10^3 is a smaller number than 4^3 .

Explain your answer.

- (b) One of the numbers below has the same value as 3.6×10^4 . Write down the number.

$$36^3 \quad 36^4 \quad (3.6 \times 10)^4 \quad 0.36 \times 10^3 \quad 0.36 \times 10^5$$

- (c) One of the numbers below has the same value as 2.5×10^{-3} . Write down the number.

$$25 \times 10^{-4} \quad 2.5 \times 10^3 \quad -2.5 \times 10^3 \quad 0.00025 \quad 2500$$

- (d) $(2 \times 10^2) \times (2 \times 10^2)$ can be written more simply as 4×10^4 .

Write the following values as simply as possible:

(i) $(3 \times 10^2) \times (2 \times 10^{-2})$

(ii) $\frac{6 \times 10^8}{2 \times 10^4}$

(KS3/98/Ma/Tier 6-8/P1)

3.5 Fractional Indices

Indices that are fractions are used to represent square roots, cube roots and other roots of numbers.

$$a^{\frac{1}{2}} = \sqrt{a} \quad \text{for example,} \quad 9^{\frac{1}{2}} = 3$$

$$a^{\frac{1}{3}} = \sqrt[3]{a} \quad \text{for example,} \quad 8^{\frac{1}{3}} = 2$$

$$a^{\frac{1}{4}} = \sqrt[4]{a} \quad \text{for example,} \quad 625^{\frac{1}{4}} = 5$$

$$a^{\frac{1}{n}} = \sqrt[n]{a}$$



Example 1

Calculate:

(a) $81^{\frac{1}{2}}$

(b) $1000^{\frac{1}{3}}$

(c) $4^{-\frac{1}{2}}$



Solution

$$\begin{aligned} \text{(a)} \quad 81^{\frac{1}{2}} &= \sqrt{81} \\ &= 9 \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad 1000^{\frac{1}{3}} &= \sqrt[3]{1000} \\ &= 10 \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad 4^{-\frac{1}{2}} &= \frac{1}{4^{\frac{1}{2}}} \\ &= \frac{1}{\sqrt{4}} \\ &= \frac{1}{2} \end{aligned}$$



Exercises

1. Calculate:

(a) $49^{\frac{1}{2}}$

(b) $64^{\frac{1}{2}}$

(c) $16^{\frac{1}{2}}$

(d) $81^{-\frac{1}{2}}$

(e) $100^{-\frac{1}{2}}$

(f) $25^{-\frac{1}{2}}$

(g) $9^{\frac{1}{2}}$

(h) $36^{-\frac{1}{2}}$

(i) $144^{\frac{1}{2}}$

2. Calculate:

(a) $8^{\frac{1}{3}}$

(b) $8^{-\frac{1}{3}}$

(c) $125^{\frac{1}{3}}$

(d) $64^{-\frac{1}{3}}$

(e) $216^{\frac{1}{3}}$

(f) $1000000^{-\frac{1}{3}}$

3. Calculate:

(a) $32^{\frac{1}{5}}$

(b) $64^{-\frac{1}{2}}$

(c) $10000^{\frac{1}{4}}$

(d) $81^{-\frac{1}{4}}$

(e) $625^{\frac{1}{4}}$

(f) $100000^{-\frac{1}{5}}$

4. Calculate:

(a) $\left(\frac{4 \times 8}{2}\right)^{\frac{1}{2}}$

(b) $\left(\frac{9 \times 27}{3}\right)^{\frac{1}{4}}$

(c) $\left(\frac{125 \times 5}{25}\right)^{\frac{1}{2}}$

(d) $\left(\frac{625}{5}\right)^{-\frac{1}{3}}$

5. Is each of the following statements *true* or *false*:

(a) $16^{\frac{1}{2}} = 8$

(b) $16^{\frac{1}{4}} = 2$

(c) $81^{\frac{1}{3}} = 9$

(d) $\left(\frac{1}{100}\right)^{-\frac{1}{2}} = 10$

6. Simplify:

(a) $(x^9)^{\frac{1}{3}}$

(b) $(a^{10})^{-\frac{1}{2}}$

(c) $\frac{a}{a^{\frac{1}{2}}}$

(d) $\frac{a^{\frac{1}{2}}}{a}$

7. Simplify:

(a) $\frac{x^{\frac{3}{2}}}{x}$

(b) $\frac{x}{x^{\frac{2}{3}}}$

(c) $\frac{a^{\frac{1}{3}}}{a}$

(d) $\frac{a^{\frac{1}{3}}}{a^{\frac{1}{2}}}$

8. Calculate:

(a) $4^{-\frac{1}{2}} + 4^{\frac{1}{2}}$

(b) $\left(9^0 + 9^{\frac{1}{2}}\right)^{\frac{1}{2}}$

(c) $\left(256^{\frac{1}{2}}\right)^{\frac{1}{2}}$

(d) $(9 - 9^0)^{\frac{1}{3}}$