

Pacific Journal of Mathematics

METHODS OF SUMMATION

GORDON MARSHALL PETERSEN

METHODS OF SUMMATION

G. M. PETERSEN

1. Methods of Rogosinski and Bernstein. In this note we shall discuss certain matrix methods of summation, though otherwise §1 and §2 are unrelated. In this section we wish to consider some of the properties of the method (B^h) , where we say that a series $\sum_{\nu=0}^{\infty} u_{\nu}$ is summable (B^h) when

$$B_n^h = \sum_{\nu=0}^n u_{\nu} \cos \frac{\pi}{2} \left(\frac{\nu}{n+h} \right) \rightarrow S, \quad n \rightarrow \infty.$$

The method (B^h) has been the subject of recent papers by Agnew [1], Karamata [5, 6], and Petersen [7]. It has been shown in the papers by Agnew and Petersen that for $h > 1/2$ the method (B^h) is equivalent to the arithmetic means of Cesaro (C), and in the paper by Agnew that for $0 < h < 1/2$ the method is equivalent to methods stronger than (C).

We shall now construct examples after a method of Hurwitz [4], to show that for $h < 0$ the method (B^h) sums a series not summable (C). Hence, since all series summable (C) are summable (B^h) , we shall have proved that (B^h) is stronger than (C).

We shall first consider $-1 < h < 0$, so that all the coefficients in any row are positive except the n th coefficient $\cos \{ \pi n / [2(n+h)] \}$. We choose $u_0 > 1$ and assume that the first $m-1$ terms of the series $\sum_{\nu=0}^{\infty} u_{\nu}$ are known. Then we select u_m so that

$$B_m^h = \sum_{\nu=0}^m u_{\nu} \cos \frac{\pi}{2} \left(\frac{\nu}{m+h} \right) = 0,$$

or

$$- u_m \cos \frac{\pi}{2} \left(\frac{m}{m+h} \right) = \sum_{\nu=0}^{m-1} u_{\nu} \cos \frac{\pi}{2} \left(\frac{\nu}{m+h} \right).$$

Received December 31, 1952.

Pacific J. Math. 4 (1954), 73-77

All of the u_ν are positive; and since

$$\frac{u_m}{u_{m-2}} \geq \frac{\sin \frac{\pi}{2} \left(\frac{2+h}{m+h} \right)}{-\sin \frac{\pi}{2} \left(\frac{h}{m+h} \right)} \simeq -\left(\frac{2}{h} + 1 \right)$$

for $-1 < h < 0$, the u_ν do not satisfy $u_n = o(n)$, and hence $\sum_{\nu=0}^{\infty} u_\nu$ is not summable (C); see [3].

If $h \leq -1$, we consider

$$B_m^h = \sum_{\nu=0}^{m-1} \left[\cos \frac{\pi}{2} \left(\frac{\nu}{m+h} \right) - \cos \frac{\pi}{2} \left(\frac{\nu+1}{m+h} \right) \right] S_\nu + \cos \frac{\pi}{2} \left(\frac{m}{m+h} \right) S_m.$$

Here again we select positive increasing S_ν so that $B_\nu^h = 0$ for $\nu \leq m-1$. Under the assumption that $S_\nu \geq \nu$, $\nu \leq m-1$, we shall show that $S_m \geq m$. Observing that the first $m-1$ coefficients of the S_ν are positive, we have (setting $\pi/[2(m+h)] = \theta$):

$$\begin{aligned} -\cos m\theta &\geq \sum_{\nu=0}^{m-1} [\cos \nu\theta - \cos (\nu+1)\theta] \nu \\ &= \sum_{\nu=0}^{m-1} \cos \nu\theta - (m-1) \cos m\theta \\ &= \Re \sum_{\nu=0}^{m-1} e^{i\nu\theta} - (m-1) \cos m\theta \\ &= \Re \frac{1 - e^{im\theta}}{1 - e^{i\theta}} - (m-1) \cos m\theta \\ &= \Re \frac{i(e^{-(i\theta)/2} - e^{i(m-1/2)\theta})}{2 \sin \theta/2} - (m-1) \cos m\theta \\ &\geq \left(\frac{1}{2} - \frac{\pi}{2} h \right); \end{aligned}$$

therefore,

$$S_m \geq \left(\frac{1}{2} - \frac{\pi}{2} h \right) \frac{m+h}{-h} \times \frac{2}{\pi} \geq qm, \quad q > 1.$$

Hence the series constructed does not satisfy the condition $S_n = o(n)$, and is not summable (C).

2. A Nörlund method. The method defined by

$$\sigma_n = \left(1 - \frac{1}{n+3} \right) S_n + \frac{1}{n+3} S_{n+1}$$

has been used as an example in a recent paper by Agnew [2]. We shall treat this method in a manner similar to that in which the method

$$t_n = (1-\alpha)S_{n-1} + \alpha S_n$$

is treated in [7].

THEOREM. *If*

$$\sigma_n = \left[\left(1 - \frac{1}{n+3} \right) S_n + \frac{1}{n+3} S_{n+1} \right] \rightarrow \sigma,$$

then

$$S_n = C \cdot (-1)^{n-1} (n+1)! + \sigma'_n,$$

where σ'_n is convergent to σ and C is a constant.

Proof. Since (we may assume $S_0 = 0$)

$$(n+2) \sigma_{n-1} = (n+1) S_{n-1} + S_n$$

$$(n+1) \sigma_{n-2} = n S_{n-2} + S_{n-1}$$

$$\cdot \quad \cdot \quad \cdot$$

$$\cdot \quad \cdot \quad \cdot$$

$$\cdot \quad \cdot \quad \cdot$$

$$3 \sigma_0 = 2 S_0 + S_1 ,$$

we have

$$S_n = (n+2)\sigma_{n-1} - (n+1)^2\sigma_{n-2} + n^2(n+1)\sigma_{n-3}$$

$$- (n-1)^2 n(n+1)\sigma_{n-4} + \cdots + (-1)^{n-2} 3^2 \cdot 4 \cdot 5 \cdot 6 \cdots (n+1)\sigma_0,$$

or

$$\begin{aligned} S_n = & (-1)^{n-1}(n+1)! \left[(-1)^{n-1} \frac{n+2}{(n+1)!} \sigma_{n-1} + (-1)^{n-2} \frac{n+1}{n!} \sigma_{n-2} + \cdots \right. \\ & \left. + (-1)^\nu \frac{\nu+3}{(\nu+2)!} \sigma_\nu + \cdots + \frac{3}{2} \sigma_0 \right]. \end{aligned}$$

Let

$$(-1)^\nu \frac{\nu+3}{(\nu+2)!} \sigma_\nu = t_\nu;$$

since $\sum_{\nu=0}^{\infty} t_\nu$ is absolutely convergent ($\sigma_\nu \rightarrow \sigma$), we may write

$$\begin{aligned} t_0 + t_1 + \cdots + t_{n-1} &= C - (t_n + t_{n+1} + \cdots) \\ &= C - \frac{1}{(n+1)!} \left[\frac{n+3}{n+2} \frac{(n+2)!}{n+3} t_n + \frac{n+4}{(n+2)(n+3)} \frac{(n+3)!}{n+4} t_{n+1} + \cdots \right] \\ &= C - \frac{(-1)^n}{(n+1)!} \left[\frac{n+3}{n+2} \sigma_n - \frac{n+4}{(n+2)(n+3)} \sigma_{n+1} + \cdots \right]. \end{aligned}$$

Then

$$\begin{aligned} S_n &= (-1)^{n-1}(n+1)! [t_0 + t_1 + \cdots + t_{n-1}] \\ &= (-1)^{n-1} \cdot C \cdot (n+1)! + \left[\frac{n+3}{n+2} \sigma_n - \frac{n+4}{(n+2)(n+3)} \sigma_{n+1} + \cdots \right] \\ &= (-1)^{n-1} \cdot C \cdot (n+1)! + \frac{n+3}{n+2} \sigma_n \\ &\quad - \frac{1}{n+2} \left[\frac{n+4}{n+3} \sigma_{n+1} - \frac{n+5}{(n+3)(n+4)} \sigma_{n+2} + \cdots \right] \\ &= (-1)^{n-1} \cdot C \cdot (n+1)! + \frac{n+3}{n+2} \sigma_n - \frac{1}{n+2} O(1) \end{aligned}$$

$$= (-1)^{n+1} \cdot C \cdot (n+1)! + \sigma_n + o(1).$$

This proves our assertion.

Obvious extensions can be made to the methods

$$\sigma_n = \left[\left(1 - \frac{1}{n+k} \right) S_n + \frac{1}{n+k} S_{n+1} \right],$$

or to iterations of these methods.

REFERENCES

1. R. P. Agnew, *Rogosinski-Bernstein trigonometric summability methods and modified arithmetic means*, Ann. of Math. (2) 56 (1952), 537-59.
2. ———, *Equivalence of methods for evaluation of sequences*, Proc. Amer. Math. Soc. 3 (1952), 550-56.
3. G. H. Hardy, *Divergent series*, Oxford, 1949.
4. W. A. Hurwitz, *Totally regular methods of summation*, Proc. London Math. Soc. (2) 26, (1926), 231-48.
5. J. Karamata, *Sur la sommabilité de S. Bernstein et quelques procédés de sommation qui s'y rattachent*, Mat. Sobornik, N.S. 21 (63) (1947), 13-24.
6. J. Karamata, *Über die Beziehung zwischen dem Bernsteinschen und Cesàroschen Limitierungsverfahren*, Math. Z. 52 (1949), 305.
7. G. M. Petersen, *A note on divergent series*, Canadian J. Math. 4 (1952), 445-54.

UNIVERSITY OF ARIZONA

UNIVERSITY OF OKLAHOMA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

M. M. SCHIFFER*

Stanford University
Stanford, California

R. P. DILWORTH

California Institute of Technology
Pasadena 4, California

E. HEWITT

University of Washington
Seattle 5, Washington

E. F. BECKENBACH**

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN

HERBERT FEDERER
MARSHALL HALL

P. R. HALMOS

HEINZ HOPF
R. D. JAMES

BØRGE JESSEN

PAUL LÉVY
GEORGE POLYA

J. J. STOKER

E. G. STRAUS
KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA, BERKELEY

UNIVERSITY OF CALIFORNIA, DAVIS

UNIVERSITY OF CALIFORNIA, LOS ANGELES

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD RESEARCH INSTITUTE

STANFORD UNIVERSITY

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

*

AMERICAN MATHEMATICAL SOCIETY

HUGHES AIRCRAFT COMPANY

SHELL DEVELOPMENT COMPANY

* To be succeeded in 1955, by H. L. Royden, Stanford University, Stanford, California.

** To be succeeded in 1955, by E. G. Straus, University of California, Los Angeles 24, Calif.

Vari-Type Composition by

Elaine Barth

Printed in the United States of America by
Edwards Brothers, Inc., Ann Arbor, Michigan

UNIVERSITY OF CALIFORNIA PRESS * BERKELEY AND LOS ANGELES
COPYRIGHT 1954 BY PACIFIC JOURNAL OF MATHEMATICS

Pacific Journal of Mathematics

Vol. 4, No. 1

May, 1954

Hugh D. Brunk, <i>On the growth of functions having poles or zeros on the positive real axis</i>	1
J. Copping, <i>Application of a theorem of Pólya to the solution of an infinite matrix equation</i>	21
James Richard Jackson, <i>On the existence problem of linear programming</i>	29
Victor Klee, <i>Invariant extension of linear functionals</i>	37
Shu-Teh Chen Moy, <i>Characterizations of conditional expectation as a transformation on function spaces</i>	47
Hukukane Nikaidô, <i>On von Neumann's minimax theorem</i>	65
Gordon Marshall Petersen, <i>Methods of summation</i>	73
G. Power, <i>Some perturbed electrostatic fields</i>	79
Murray Harold Protter, <i>The two noncharacteristic problem with data partly on the parabolic line</i>	99
S. E. Rauch, <i>Mapping properties of Cesàro sums of order two of the geometric series</i>	109
Gerson B. Robison, <i>Invariant integrals over a class of Banach spaces</i>	123
Richard Steven Varga, <i>Eigenvalues of circulant matrices</i>	151