



**UCF COASTAL**  
NATIONAL CENTER FOR INTEGRATED COASTAL RESEARCH

# COMPOUND FLOODING: EXAMPLES, METHODS, AND CHALLENGES



***Thomas Wahl***

Civil, Environmental and Construction  
Engineering & National Center for  
Integrated Coastal Research  
University of Central Florida

# Definition(s) of compound events

## IPCC, 2012:

*(1) two or more extreme events occurring simultaneously or successively; (2) combinations of extreme events with underlying conditions that amplify the impact of the events; or (3) combinations of events that are not themselves extremes but lead to an extreme event or impact when combined. The contributing events can be of similar (clustered multiple events) or different type(s).*


## Leonard et al., 2014:

*A compound event is an extreme impact that depends on multiple statistically dependent variables or events.*

## Zscheischler et al., 2018:

*The combination of multiple drivers and/or hazards that contributes to societal or environmental impacts.*


# Definition(s) of compound events




*Zscheischler et al., 2018*

# Types of compound flooding

## Coastal



## River confluences



*Bender et al., 2016*

# Examples

Compound flooding in the upper Danube basin (city of Passau) from high discharges into the confluence of the rivers Danube, Inn and Ilz.



Compound flooding in the city of Jacksonville from the combination of storm surge and high discharge of the St. John's river during hurricane Irma.



Compound flooding in Galveston from extreme precipitation and moderate but long-lasting storm surge during hurricane Harvey.






Yarmouth, 27 Oct 2013



Whitby, 5 Dec 2013



Chesil Jan 2014



Aberystwyth, Feb 2014



Dawlish, Feb 2013


27-28 Oct, 'St Jude' Storm  
99 mph winds  
Yarmouth: floods neap tide

5-6 Dec, Xaver Storm  
East coast surge  
Repeat of 1953?

1-3 Jan,  
Wales, Dorset  
badly hit

1- 5 & 14 Feb,  
Destruction Dawlish  
railway

3 Mar,  
flooding in  
Jersey



2013

2014

Photo source: BBC

# Methods

Dependence of marginal parameters

statistical independence  
(e.g.  $\chi^2$ -Test)

multiplication of marginal  
exceedance probabilities

statistical dependence  
(e.g.  $r \neq 0$ )

multivariate models

Marginal distributions




same marginal distributions

mixed marginal distributions

multivariate parametric models  
(e.g. bivariate Gumbel model)


Copula models  
(e.g. Gumbel, Frank, Clayton)

Archimedean  
Copula family



# Non-stationarity

Non-stationarity in the multivariate case can come from either of the relevant variables (e.g. trends, cycles etc.) or changes in their dependency.



**Green:** hypothetical distribution of two climatic drivers in the present climate.


**Blue:** a future climate with shift in mean, variability and correlation between the drivers.

**Purple:** future climate with an increase in dependence in the upper tail of both drivers.

# Non-stationarity

The risk of compound flooding is assessed using sea level records from 30 tide gauges (> 30 years of data) and precipitation data from stations within 25 km (averaged into indices for each tide gauge).


The dependency between the two proxies and the spatial and temporal variability are analyzed:



# Non-stationarity

The risk of compound flooding is assessed using sea level records from 30 tide gauges (> 30 years of data) and precipitation data from stations within 25 km (averaged into indices for each tide gauge).

The dependency between the two proxies and the spatial and temporal variability are analyzed:




# Non-stationarity

## Extreme Value Analysis:

We select a random extreme event combination (comprised of 120 mm daily rainfall and 115 cm storm surge; i.e. similar to tropical storm Doria) and quantify the effect of the increased dependency on multivariate (“AND”) return periods:

- Independence assumption:  $RP_{\text{mult, ind}} \approx 245 \text{ yrs}$
- Using small correlation as observed in the 1940s:  $RP_{\text{mult, } \tau_{1940s}} \approx 105 \text{ yrs}$
- Using large correlation as observed recently:  $RP_{\text{mult, } \tau_{2000s}} \approx 42 \text{ yrs}$



# The way forward

- Identify additional key variable and event combinations needing scrutiny. Use bottom-up approaches and perform system stress tests to identify vulnerabilities.
- Use appropriate statistical methods to simulate dependence in time (i.e., temporal clustering), space (i.e., spatial footprints) and across multiple variables.
- Identify data and model requirements for documenting, understanding, simulating and attributing compound events.
- Incorporating compound events into impact assessments and disaster risk mitigation works.

**Requires close collaboration and communication between scientists from various fields (natural sciences, engineering, social sciences) as well as stakeholders and policy makers.**



**Thank you for your  
attention!**