



## **AP<sup>®</sup> Physics C: Electricity and Magnetism 2012 Scoring Guidelines**

### **The College Board**

The College Board is a mission-driven not-for-profit organization that connects students to college success and opportunity. Founded in 1900, the College Board was created to expand access to higher education. Today, the membership association is made up of more than 5,900 of the world's leading educational institutions and is dedicated to promoting excellence and equity in education. Each year, the College Board helps more than seven million students prepare for a successful transition to college through programs and services in college readiness and college success — including the SAT<sup>®</sup> and the Advanced Placement Program<sup>®</sup>. The organization also serves the education community through research and advocacy on behalf of students, educators, and schools. The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities, and concerns.

© 2012 The College Board. College Board, Advanced Placement Program, AP, SAT and the acorn logo are registered trademarks of the College Board. All other products and services may be trademarks of their respective owners. Permission to use copyrighted College Board materials may be requested online at: [www.collegeboard.com/inquiry/cbpermit.html](http://www.collegeboard.com/inquiry/cbpermit.html).

Visit the College Board on the Web: [www.collegeboard.org](http://www.collegeboard.org).

AP Central is the official online home for the AP Program: [apcentral.collegeboard.org](http://apcentral.collegeboard.org).



# AP<sup>®</sup> PHYSICS

## 2012 SCORING GUIDELINES

### General Notes About 2012 AP Physics Scoring Guidelines

1. The solutions contain the most common method of solving the free-response questions and the allocation of points for this solution. Some also contain a common alternate solution. Other methods of solution also receive appropriate credit for correct work.
2. Generally, double penalty for errors is avoided. For example, if an incorrect answer to part (a) is correctly substituted into an otherwise correct solution to part (b), full credit will usually be awarded in part (b). One exception to this practice may occur in cases where the numerical answer to a later part should easily be recognized as wrong, for example, a speed faster than the speed of light in vacuum.
3. Implicit statements of concepts normally receive credit. For example, if the use of an equation expressing a particular concept is worth 1 point, and a student's solution contains the application of that equation to the problem but the student does not write the basic equation, the point is still awarded. However, when students are asked to derive an expression, it is normally expected that they will begin by writing one or more fundamental equations, such as those given on the AP Physics Exam equation sheets. For a description of the use of such terms as "derive" and "calculate" on the exams, and what is expected for each, see "The Free-Response Sections — Student Presentation" in the *AP Physics Course Description*.
4. The scoring guidelines typically show numerical results using the value  $g = 9.8 \text{ m/s}^2$ , but use of  $10 \text{ m/s}^2$  is of course also acceptable. Solutions usually show numerical answers using both values when they are significantly different.
5. Strict rules regarding significant digits are usually not applied to numerical answers. However, in some cases answers containing too many digits may be penalized. In general, two to four significant digits are acceptable. Numerical answers that differ from the published answer owing to differences in rounding throughout the question typically receive full credit. Exceptions to these guidelines usually occur when rounding makes a difference in obtaining a reasonable answer. For example, suppose a solution requires subtracting two numbers that should have five significant figures and that differ starting with the fourth digit (e.g., 20.295 and 20.278). Rounding to three digits will eliminate the level of accuracy required to determine the difference in the numbers, and some credit may be lost.

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2012 SCORING GUIDELINES**

**Question 1**

**15 points total**

**Distribution  
of points**

(a) 3 points

For an expression of Gauss's Law

1 point

$$\frac{Q_i}{\epsilon_0} = \oint \mathbf{E} \cdot d\mathbf{A}$$

For a correct intermediate step indicating that the area of the Gaussian surface is  $4\pi r^2$

1 point

$$\frac{Q_i}{\epsilon_0} = E(4\pi r^2)$$

For a correct final expression, specifically using  $Q_i$

1 point

$$E(r) = \frac{1}{4\pi\epsilon_0} \frac{Q_i}{r^2} \quad \text{or} \quad E(r) = \frac{kQ_i}{r^2}$$

(b) 2 points

For indicating that the enclosed charge is the sum of the inner and outer charges

1 point

$$\frac{Q_i + Q_o}{\epsilon_0} = \oint \mathbf{E} \cdot d\mathbf{A}$$

For a correct expression for the electric field

1 point

$$E(r) = \frac{1}{4\pi\epsilon_0} \frac{(Q_i + Q_o)}{r^2}$$

Note: The correct expression by itself earns both points.

(c) 2 points

For using the integral definition of potential in terms of electric field

1 point

$$V(r) - V_\infty = -\int_\infty^r \mathbf{E} \cdot d\mathbf{r}$$

$$V(r) = -\int_\infty^r \frac{1}{4\pi\epsilon_0} \frac{(Q_i + Q_o)}{r^2} dr$$

$$V(r) = -\frac{(Q_i + Q_o)}{4\pi\epsilon_0} \left[ -\frac{1}{r} \right]_\infty^r$$

For the correct expression

1 point

$$V(r) = \frac{1}{4\pi\epsilon_0} \frac{(Q_i + Q_o)}{r}$$

Note: The correct expression by itself earns both points.

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2012 SCORING GUIDELINES**

**Question 1 (continued)**


|     |  | <b>Distribution<br/>of points</b> |
|-----|--|-----------------------------------|
| (c) | continued  |                                   |
|     | <i>Alternate solution</i>  | <i>Alternate points</i>           |
|     | <i>Outside the shells, the charges on each can be treated as point charges at their centers.</i>   |                                   |
|     | <i>For using the concept of summation of point charge potentials</i>   | <i>1 point</i>                    |
|     | $V(r) = \frac{1}{4\pi\epsilon_0} \sum_j \frac{q_j}{r_j}$   |                                   |
|     | $V(r) = \frac{1}{4\pi\epsilon_0} \left( \frac{Q_i}{r} + \frac{Q_o}{r} \right)$   |                                   |
|     | <i>For the correct expression (the correct expression by itself earns both points)</i>   | <i>1 point</i>                    |
|     | $V(r) = \frac{1}{4\pi\epsilon_0} \frac{(Q_i + Q_o)}{r}$  |                                   |
| (d) | 1 point  |                                   |
|     | The answer from part (c), with $Q_T = Q_i + Q_o$ , can be solved for $Q_T$ . Values at the outer shell are then used to determine a numerical value. |                                   |
|     | For correct work resulting in the correct value, with units  | 1 point                           |
|     | $V(r) = \frac{1}{4\pi\epsilon_0} \frac{Q_T}{r}, \quad r \geq 0.20 \text{ m}$   |                                   |
|     | $Q_T = 4\pi\epsilon_0 r V(r) = \left( \frac{(0.20 \text{ m})(100 \text{ V})}{(9.0 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2)} \right)$         |                                   |
|     | $Q_T = 2.2 \times 10^{-9} \text{ C}$   |                                   |

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2012 SCORING GUIDELINES**

**Question 1 (continued)**

**Distribution  
of points**


(e) 3 points



- |   |         |
|---|---------|
| For a segment indicating an $E$ -field of 0 for $r < 0.10$ m, explicitly drawn  | 1 point |
| For a segment that is concave down and negative for $0.10$ m $< r < 0.20$ m   | 1 point |
| For a segment that is concave up and positive for $r > 0.20$ m. The line must not touch or cross the horizontal axis. | 1 point |

Note: The labels on the vertical axis are not to scale and are not required to receive full credit.

(f) 4 points



- |   |         |
|---|---------|
| For a continuous set of segments that have slope discontinuities at $r = 0.10$ m and at $r = 0.20$ m                  | 1 point |
| For a segment indicating a constant negative potential for $r < 0.10$ m   | 1 point |
| For a segment that is increasing, concave down, and crosses the $r$ axis, for $0.10$ m $< r < 0.20$ m                 | 1 point |
| For a segment that is concave up and positive for $r > 0.20$ m. The line must not touch or cross the horizontal axis. | 1 point |

Note: The labels on the vertical axis are not scored and are not required to receive full credit.


**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2012 SCORING GUIDELINES**

**Question 2**

**15 points total**

**Distribution  
of points**

(a) 4 points



For a correct label on each axis (one including resistance, one including length), that leads to a linear graph

1 point

For two linear scales, one for each axis, corresponding to the labels and occupying at least three-quarters of each axis

1 point

For reasonably correctly plotted points according to the scale

1 point

For a reasonable best-fit straight line

1 point

Note: Circles on the graph are to indicate points chosen to calculate slope in part (b). They are not necessary to receive credit.

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2012 SCORING GUIDELINES**

**Question 2 (continued)**


|       |   | <b>Distribution<br/>of points</b> |
|-------|---|-----------------------------------|
| (b)   | 3 points  |                                   |
|       | For correctly using the equation $R = \rho L/A$ to solve for the resistivity<br>$\rho = (R/L)A$   | 1 point                           |
|       | For calculating the slope $m$ from two points that lie on the indicated best-fit line<br><u>Example (using the two points indicated on the graph)</u><br>$m = \Delta R/\Delta L = (340 \times 10^3 \Omega - 140 \times 10^3 \Omega)/(0.076 \text{ m} - 0.032 \text{ m})$ $m = 4.55 \times 10^6 \Omega/\text{m}$                         | 1 point                           |
|       | <u>Note:</u> The slope point can also be earned for indicating that a linear regression function on a calculator was used (exact answer required): $m = 4.5 \times 10^6 \Omega/\text{m}$ (to two significant figures).<br>$\rho = mA = mtw$<br>$\rho = (4.5 \times 10^6 \Omega/\text{m})(1.0 \times 10^{-4} \text{ m})(0.02 \text{ m})$ |                                   |
|       | For a correct answer, with supporting work<br>$\rho = 9.0 \Omega \cdot \text{m}$ ( $\pm 0.2 \Omega \cdot \text{m}$ is acceptable)   | 1 point                           |
| (c)   | 3 points  |                                   |
|       | For correctly stating the RC circuit time constant equation<br>$\tau = RC$  | 1 point                           |
|       | For the correct value of the equivalent resistance<br>$\frac{1}{R_p} = \frac{1}{R_4} + \frac{1}{R_5} = \frac{1}{(370 \text{ k}\Omega)} + \frac{1}{(440 \text{ k}\Omega)}$ $R_p = 201 \text{ k}\Omega$   | 1 point                           |
|       | For the correct value of the time constant (units not evaluated here)<br>$\tau = R_p C = (201 \text{ k}\Omega)(10 \mu\text{F})$<br>$\tau = 2.01 \text{ s}$  | 1 point                           |
| Units | 1 point   |                                   |
|       | For correct units for the answers in both parts (b) and (c)   | 1 point                           |

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2012 SCORING GUIDELINES**

**Question 2 (continued)**

**Distribution  
of points**

(d) 4 points



- |   |         |
|---|---------|
| For correctly indicating the battery voltage as an asymptote for the $V_C$ curve (or in the absence of a correct $V_C$ curve, as the maximum of the resistor curves)                            | 1 point |
| For a capacitor voltage curve starting at the origin, increasing, concave down, and appropriately labeled   | 1 point |
| For a resistor voltage curve starting at the intersection of the asymptotic voltage with the voltage axis, decreasing and concave up (asymptotic with the time axis), and appropriately labeled | 1 point |
| For any indication that $V_{R4} = V_{R5}$   | 1 point |



**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2012 SCORING GUIDELINES**

**Question 3**

**15 points total**

**Distribution  
of points**

(a) 1 point

$$\phi_m = \int \mathbf{B} \cdot d\mathbf{A}$$

For a correct answer

$$\phi_m = B_0 L h_0$$

1 point

(b) 2 points



For a correct direction arrow

1 point

For a valid justification using Lenz's Law or the right-hand rule, if the direction is also correct

1 point

Examples

The flux is decreasing as the area is decreasing, and a current to the right would cause an inward magnetic field that would increase flux.

The force on the falling positive charge carriers is to the right, which causes a conventional current to the right.

(c) 3 points

For a correct relationship between current, voltage, and resistance

1 point

$$I = \frac{\mathcal{E}}{R}$$

For a correct relationship between the induced emf and the magnetic field

1 point

$$|\mathcal{E}| = \frac{d\phi_m}{dt}$$

$$|\mathcal{E}| = B_0 L \frac{dh}{dt}$$

$$|\mathcal{E}| = B_0 L v$$

For a correct answer

1 point

$$I = B_0 L v / R$$

Note: If only the correct answer is given, with no accompanying work, only 1 point can be awarded.

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2012 SCORING GUIDELINES**

**Question 3 (continued)**

|     |   | <b>Distribution<br/>of points</b> |
|-----|---|-----------------------------------|
| (d) | 4 points  |                                   |
|     | For a correct net force equation showing opposite directions for the gravitational and magnetic forces, $F_g$ and $F_M$ | 1 point                           |
|     | $\sum F = ma = F_g - F_M = mg - F_M$ $a = g - \frac{F_M}{m}$  |                                   |
|     | For using an appropriate equation to find $F_M$   | 1 point                           |
|     | $F_M = \int I  d\boldsymbol{\ell} \times \mathbf{B}  = ILB_0$   |                                   |
|     | For substituting the current from part (c)  | 1 point                           |
|     | $F_M = \left( \frac{B_0Lv}{R} \right) LB_0 = \frac{B_0^2L^2v}{R}$   |                                   |
|     | For expressing acceleration $a$ as $dv/dt$  | 1 point                           |
|     | $\frac{dv}{dt} = g - \frac{B_0^2L^2v}{mR}$  |                                   |
| (e) | 2 points  |                                   |
|     | For setting the gravitational force equal to the magnetic force<br>$a = 0$ ; therefore $F_M = F_g$                      | 1 point                           |
|     | For correct substitution of expressions for the forces  | 1 point                           |
|     | $mg = ILB_0 = B_0^2L^2v_T/R$ $v_T = \frac{mgR}{B_0^2L^2}$   |                                   |
|     | <u>Note:</u> If the correct expression for $v_T$ is stated without support, 2 points are awarded.                       |                                   |

**AP<sup>®</sup> PHYSICS C: ELECTRICITY AND MAGNETISM  
2012 SCORING GUIDELINES**

**Question 3 (continued)**

|     |  | <b>Distribution<br/>of points</b> |
|-----|--|-----------------------------------|
| (f) | 3 points   |                                   |
|     | For correctly checking “Increases”   | 1 point                           |
|     | <u>Note:</u> If an incorrect choice is made for the change in terminal speed, the justification points cannot be earned.   |                                   |
|     | For indicating the inverse relationship between resistance and current   | 1 point                           |
|     | For indicating that a smaller current produces a smaller magnetic force on the bar, leading to the conclusion that to achieve a magnetic force equal to the bar’s weight, the bar must be moving faster to produce the necessary current in the bar. | 1 point                           |
|     | <u>Note:</u> If the only justification is stating that $v_T \propto R$ from the equation for $v_T$ in part (e), only 1 justification point is awarded, because the question specifically asks for the answer in terms of forces on the crossbar.     |                                   |