

2019

AP[®]

 CollegeBoard

AP[®] Physics 2: Algebra-Based Free-Response Questions

© 2019 The College Board. College Board, Advanced Placement, AP, AP Central, and the acorn logo are registered trademarks of the College Board. Visit the College Board on the web: collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

AP[®] PHYSICS 2 TABLE OF INFORMATION

CONSTANTS AND CONVERSION FACTORS	
Proton mass, $m_p = 1.67 \times 10^{-27}$ kg Neutron mass, $m_n = 1.67 \times 10^{-27}$ kg Electron mass, $m_e = 9.11 \times 10^{-31}$ kg Avogadro's number, $N_0 = 6.02 \times 10^{23}$ mol ⁻¹ Universal gas constant, $R = 8.31$ J/(mol·K) Boltzmann's constant, $k_B = 1.38 \times 10^{-23}$ J/K	Electron charge magnitude, $e = 1.60 \times 10^{-19}$ C 1 electron volt, $1 \text{ eV} = 1.60 \times 10^{-19}$ J Speed of light, $c = 3.00 \times 10^8$ m/s Universal gravitational constant, $G = 6.67 \times 10^{-11}$ m ³ /kg·s ² Acceleration due to gravity at Earth's surface, $g = 9.8$ m/s ²
1 unified atomic mass unit, Planck's constant, Vacuum permittivity, Coulomb's law constant, $k = 1/4\pi\epsilon_0 = 9.0 \times 10^9$ N·m ² /C ² Vacuum permeability, Magnetic constant, $k' = \mu_0/4\pi = 1 \times 10^{-7}$ (T·m)/A 1 atmosphere pressure,	$1 \text{ u} = 1.66 \times 10^{-27}$ kg = 931 MeV/c ² $h = 6.63 \times 10^{-34}$ J·s = 4.14 × 10 ⁻¹⁵ eV·s $hc = 1.99 \times 10^{-25}$ J·m = 1.24 × 10 ³ eV·nm $\epsilon_0 = 8.85 \times 10^{-12}$ C ² /N·m ² $\mu_0 = 4\pi \times 10^{-7}$ (T·m)/A $1 \text{ atm} = 1.0 \times 10^5$ N/m ² = 1.0 × 10 ⁵ Pa

UNIT SYMBOLS	meter, m	mole, mol	watt, W	farad, F
	kilogram, kg	hertz, Hz	coulomb, C	tesla, T
	second, s	newton, N	volt, V	degree Celsius, °C
	ampere, A	pascal, Pa	ohm, Ω	electron volt, eV
	kelvin, K	joule, J	henry, H	

PREFIXES		
Factor	Prefix	Symbol
10 ¹²	tera	T
10 ⁹	giga	G
10 ⁶	mega	M
10 ³	kilo	k
10 ⁻²	centi	c
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

VALUES OF TRIGONOMETRIC FUNCTIONS FOR COMMON ANGLES							
θ	0°	30°	37°	45°	53°	60°	90°
$\sin \theta$	0	1/2	3/5	$\sqrt{2}/2$	4/5	$\sqrt{3}/2$	1
$\cos \theta$	1	$\sqrt{3}/2$	4/5	$\sqrt{2}/2$	3/5	1/2	0
$\tan \theta$	0	$\sqrt{3}/3$	3/4	1	4/3	$\sqrt{3}$	∞

The following conventions are used in this exam.

- I. The frame of reference of any problem is assumed to be inertial unless otherwise stated.
- II. In all situations, positive work is defined as work done on a system.
- III. The direction of current is conventional current: the direction in which positive charge would drift.
- IV. Assume all batteries and meters are ideal unless otherwise stated.
- V. Assume edge effects for the electric field of a parallel plate capacitor unless otherwise stated.
- VI. For any isolated electrically charged object, the electric potential is defined as zero at infinite distance from the charged object

AP[®] PHYSICS 2 EQUATIONS

MECHANICS	ELECTRICITY AND MAGNETISM
$v_x = v_{x0} + a_x t$	$ \vec{F}_E = \frac{1}{4\pi\epsilon_0} \frac{ q_1 q_2 }{r^2}$
$x = x_0 + v_{x0} t + \frac{1}{2} a_x t^2$	$\vec{E} = \frac{\vec{F}_E}{q}$
$v_x^2 = v_{x0}^2 + 2a_x(x - x_0)$	$ \vec{E} = \frac{1}{4\pi\epsilon_0} \frac{ q }{r^2}$
$\vec{a} = \frac{\sum \vec{F}}{m} = \frac{\vec{F}_{net}}{m}$	$\Delta U_E = q\Delta V$
$ \vec{F}_f \leq \mu \vec{F}_n $	$V = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$
$a_c = \frac{v^2}{r}$	$ \vec{E} = \left \frac{\Delta V}{\Delta r} \right $
$\vec{p} = m\vec{v}$	$\Delta V = \frac{Q}{C}$
$\Delta \vec{p} = \vec{F} \Delta t$	$C = \kappa \epsilon_0 \frac{A}{d}$
$K = \frac{1}{2} m v^2$	$E = \frac{Q}{\epsilon_0 A}$
$\Delta E = W = F_{\parallel} d = F d \cos \theta$	$U_C = \frac{1}{2} Q \Delta V = \frac{1}{2} C (\Delta V)^2$
$P = \frac{\Delta E}{\Delta t}$	$I = \frac{\Delta Q}{\Delta t}$
$\theta = \theta_0 + \omega_0 t + \frac{1}{2} \alpha t^2$	$R = \frac{\rho \ell}{A}$
$\omega = \omega_0 + \alpha t$	$P = I \Delta V$
$x = A \cos(\omega t) = A \cos(2\pi f t)$	$I = \frac{\Delta V}{R}$
$x_{cm} = \frac{\sum m_i x_i}{\sum m_i}$	$R_s = \sum_i R_i$
$\vec{a} = \frac{\sum \vec{\tau}}{I} = \frac{\vec{\tau}_{net}}{I}$	$\frac{1}{R_p} = \sum_i \frac{1}{R_i}$
$\tau = r_{\perp} F = r F \sin \theta$	$C_p = \sum_i C_i$
$L = I \omega$	$\frac{1}{C_s} = \sum_i \frac{1}{C_i}$
$\Delta L = \tau \Delta t$	$B = \frac{\mu_0 I}{2\pi r}$
$K = \frac{1}{2} I \omega^2$	$\vec{F}_M = q\vec{v} \times \vec{B}$
$ \vec{F}_s = k \vec{x} $	$ \vec{F}_M = q\vec{v} \sin \theta \vec{B} $
$a = \text{acceleration}$	$\vec{F}_M = I \vec{\ell} \times \vec{B}$
$A = \text{amplitude}$	$ \vec{F}_M = I \vec{\ell} \sin \theta \vec{B} $
$d = \text{distance}$	$\Phi_B = \vec{B} \cdot \vec{A}$
$E = \text{energy}$	$\Phi_B = \vec{B} \cos \theta \vec{A} $
$F = \text{force}$	$\mathcal{E} = -\frac{\Delta \Phi_B}{\Delta t}$
$f = \text{frequency}$	$\mathcal{E} = B \ell v$
$I = \text{rotational inertia}$	$A = \text{area}$
$K = \text{kinetic energy}$	$B = \text{magnetic field}$
$k = \text{spring constant}$	$C = \text{capacitance}$
$L = \text{angular momentum}$	$d = \text{distance}$
$\ell = \text{length}$	$E = \text{electric field}$
$m = \text{mass}$	$\mathcal{E} = \text{emf}$
$P = \text{power}$	$F = \text{force}$
$p = \text{momentum}$	$I = \text{current}$
$r = \text{radius or separation}$	$\ell = \text{length}$
$T = \text{period}$	$P = \text{power}$
$t = \text{time}$	$Q = \text{charge}$
$U = \text{potential energy}$	$q = \text{point charge}$
$v = \text{speed}$	$R = \text{resistance}$
$W = \text{work done on a system}$	$r = \text{separation}$
$x = \text{position}$	$t = \text{time}$
$y = \text{height}$	$U = \text{potential (stored) energy}$
$\alpha = \text{angular acceleration}$	$V = \text{electric potential}$
$\mu = \text{coefficient of friction}$	$v = \text{speed}$
$\theta = \text{angle}$	$\kappa = \text{dielectric constant}$
$\tau = \text{torque}$	$\rho = \text{resistivity}$
$\omega = \text{angular speed}$	$\theta = \text{angle}$
$U_s = \frac{1}{2} k x^2$	$\Phi = \text{flux}$
$\Delta U_g = mg \Delta y$	
$T = \frac{2\pi}{\omega} = \frac{1}{f}$	
$T_s = 2\pi \sqrt{\frac{m}{k}}$	
$T_p = 2\pi \sqrt{\frac{\ell}{g}}$	
$ \vec{F}_g = G \frac{m_1 m_2}{r^2}$	
$\vec{g} = \frac{\vec{F}_g}{m}$	
$U_G = -\frac{G m_1 m_2}{r}$	

AP[®] PHYSICS 2 EQUATIONS

FLUID MECHANICS AND THERMAL PHYSICS

$$\rho = \frac{m}{V}$$

$$P = \frac{F}{A}$$

$$P = P_0 + \rho gh$$

$$F_b = \rho Vg$$

$$A_1 v_1 = A_2 v_2$$

$$P_1 + \rho g y_1 + \frac{1}{2} \rho v_1^2 = P_2 + \rho g y_2 + \frac{1}{2} \rho v_2^2$$

$$\frac{Q}{\Delta t} = \frac{kA \Delta T}{L}$$

$$PV = nRT = Nk_B T$$

$$K = \frac{3}{2} k_B T$$

$$W = -P \Delta V$$

$$\Delta U = Q + W$$

A = area
F = force
h = depth
k = thermal conductivity
K = kinetic energy
L = thickness
m = mass
n = number of moles
N = number of molecules
P = pressure
Q = energy transferred to a system by heating
T = temperature
t = time
U = internal energy
V = volume
v = speed
W = work done on a system
y = height
ρ = density

MODERN PHYSICS

$$E = hf$$

$$K_{\max} = hf - \phi$$

$$\lambda = \frac{h}{p}$$

$$E = mc^2$$

E = energy
f = frequency
K = kinetic energy
m = mass
p = momentum
λ = wavelength
φ = work function

WAVES AND OPTICS

$$\lambda = \frac{v}{f}$$

$$n = \frac{c}{v}$$

$$n_1 \sin \theta_1 = n_2 \sin \theta_2$$

$$\frac{1}{s_i} + \frac{1}{s_o} = \frac{1}{f}$$

$$|M| = \left| \frac{h_i}{h_o} \right| = \left| \frac{s_i}{s_o} \right|$$

$$\Delta L = m\lambda$$

$$d \sin \theta = m\lambda$$

d = separation
f = frequency or focal length
h = height
L = distance
M = magnification
m = an integer
n = index of refraction
s = distance
v = speed
λ = wavelength
θ = angle

GEOMETRY AND TRIGONOMETRY

Rectangle
 $A = bh$

Triangle
 $A = \frac{1}{2}bh$

Circle
 $A = \pi r^2$
 $C = 2\pi r$

Rectangular solid
 $V = \ell wh$

Cylinder
 $V = \pi r^2 \ell$
 $S = 2\pi r \ell + 2\pi r^2$

Sphere
 $V = \frac{4}{3}\pi r^3$
 $S = 4\pi r^2$


A = area
C = circumference
V = volume
S = surface area
b = base
h = height
ℓ = length
w = width
r = radius

Right triangle
 $c^2 = a^2 + b^2$

$$\sin \theta = \frac{a}{c}$$

$$\cos \theta = \frac{b}{c}$$

$$\tan \theta = \frac{a}{b}$$



2019 AP[®] PHYSICS 2 FREE-RESPONSE QUESTIONS


PHYSICS 2

Section II

Time—1 hour and 30 minutes

4 Questions

Directions: Questions 1 and 4 are short free-response questions that require about 20 minutes each to answer and are worth 10 points each. Questions 2 and 3 are long free-response questions that require about 25 minutes each to answer and are worth 12 points each. Show your work for each part in the space provided after that part.




1. (10 points, suggested time 20 minutes)


The figure above shows a particle with positive charge $+Q$ traveling with a constant speed v_0 to the right and in the plane of the page. The particle is approaching a region, shown by the dashed box, that contains a constant uniform field. The effects of gravity are negligible.

(a)

- i. On the figure below, draw a possible path of the particle in the region if the region contains only an electric field directed toward the bottom of the page.




- ii. On the figure below, draw a possible path of the particle in the region if the region contains only a magnetic field directed out of the page.




- iii. For which of the previous situations is the motion more similar to that of a projectile in only a gravitational field near Earth's surface, and why?

2019 AP[®] PHYSICS 2 FREE-RESPONSE QUESTIONS




- (b) Another region of space contains an electric field directed toward the top of the page and a magnetic field directed out of the page. Both fields are constant and uniform. A horizontal beam of protons with a variety of speeds enters the region, as shown above. Protons exit the region at a variety of locations, including points 1 and 2 shown on the figure. In a coherent, paragraph-length response, explain why some protons exit the region at point 1 and others exit at point 2. Use physics principles to explain your reasoning.

2019 AP[®] PHYSICS 2 FREE-RESPONSE QUESTIONS



Circuit 1




Circuit 2

2. (12 points, suggested time 25 minutes)

The two circuits shown above contain an ideal variable power supply, an ohmic resistor of resistance R , an ammeter A , and two voltmeters V_{PS} and V_R . In circuit 1 the ammeter has negligible resistance, and in circuit 2 the ammeter has significant internal ohmic resistance r . The potential difference of the power supply is varied, and measurements of current and potential difference are recorded.

(a) The axes below can be used to graph the current measured by the ammeter as a function of the potential difference measured across the power supply. On the axes, do the following.

- Sketch a possible graph for circuit 1 and label it 1.
- Sketch a possible graph for circuit 2 and label it 2.




(b) Let ΔV_{PS} be the potential difference measured by voltmeter V_{PS} across the power supply, and let I be the current measured by the ammeter A . For each circuit, write an equation that satisfies conservation of energy, in terms of ΔV_{PS} , I , R , and r , as appropriate.

Circuit 1

Circuit 2

2019 AP[®] PHYSICS 2 FREE-RESPONSE QUESTIONS


- (c) Explain how your equations in part (b) account for any differences between graphs 1 and 2 in part (a).
- (d) In circuit 2, $R = 40\ \Omega$. When voltmeter V_{PS} reads 3.0 V, voltmeter V_R reads 2.5 V. Calculate the internal resistance r of the ammeter.
- (e) Voltmeter V_R in circuit 2 is replaced by a resistor with resistance $120\ \Omega$ to create circuit 3 shown below. Voltmeter V_{PS} still reads 3.0 V.



Circuit 3

- Calculate the equivalent resistance R_{eq} of the circuit.
- Calculate the current in each of the resistors that are in parallel.

2019 AP[®] PHYSICS 2 FREE-RESPONSE QUESTIONS



3. (12 points, suggested time 25 minutes)


A group of students use the apparatus shown above to determine the thermal conductivity of a certain type of plastic. A hot plate is used to keep water in a container boiling at a temperature of 100°C . They place a slab of the plastic with area 0.025 m^2 and thickness 0.010 m above the container so that the bottom surface of the slab is at a temperature of 100°C . They put a large block of ice with temperature 0°C on top of the plastic slab. Some of the ice melts, and the students measure the amount of water collected during a time Δt . The students correctly calculate the amount of energy Q delivered to the ice and thus determine $Q/\Delta t$. They repeat this experiment several times, each time adding an identical slab to increase the total thickness L of plastic. Their results are shown in the table below.

Energy flow rate $Q/\Delta t$ (J/s)	97	53	31	27	18
Total thickness of plastic L (m)	0.010	0.020	0.030	0.040	0.050


(a) The students want to create a graph to yield a straight line whose slope could be used to calculate the thermal conductivity of the plastic.

2019 AP[®] PHYSICS 2 FREE-RESPONSE QUESTIONS

- i. Label the axes below to indicate a pair of quantities that could be graphed to yield a straight line. Include units for the quantities.



- ii. On the grid on the previous page, create a linear graph using the values for the quantities indicated in part (a)(i). Be sure to do the following.
- Add to the data table the values of any quantities to be plotted that are not already given.
 - Scale the axes.
 - Plot the data from the table.
 - Draw a line that best represents the data.
- iii. Use the graph to calculate the thermal conductivity of the plastic.
- (b) Indicate one potential problem with the setup that could lead to an experimental value for the thermal conductivity that is different from the actual value. Use physics principles to explain the effect this problem could have on the experimental value.
- (c) The rectangle below represents a side view of the plastic slab. Draw a single arrow on the diagram representing the direction of the net flow of energy through the plastic.



2019 AP[®] PHYSICS 2 FREE-RESPONSE QUESTIONS


- (d) Describe what occurs in the plastic at the microscopic level that explains the energy flow you indicated in part (c).
- (e) An extra plastic slab sits on a wood surface, with both the plastic slab and the wood surface at room temperature. A student touches each and finds that the plastic slab feels cooler than the wood surface. Explain what causes this observation.

2019 AP[®] PHYSICS 2 FREE-RESPONSE QUESTIONS

4. (10 points, suggested time 20 minutes)

A student notices many air bubbles rising through the water in a large fish tank at an aquarium.

- (a) In the figure below, the circle represents one such air bubble, and two incoming rays of light, A and B, are shown. Ray B points toward the center of the circle. On the diagram, draw the paths of rays A and B as they go through the bubble and back into the water. Your diagram should clearly show what happens to the rays at each interface.



- (b) The bubble has a volume V_1 , the air inside it has density ρ_A , and the water around it has density ρ_W . The bubble starts at rest and has a speed v_f when it has risen a height h . Assume that the change in the bubble's volume is negligible. Derive an expression for the mechanical energy dissipated by drag forces as the bubble rises this distance. Express your answer in terms of the given quantities and fundamental constants, as appropriate.
- (c) At a particular instant, one bubble is 4.5 m below the water's surface. The surface of the water is at sea level, and the density of the water is 1000 kg/m^3 .
- Determine the absolute pressure in the bubble at this location.
 - The bubble has a volume V_1 when it is 4.5 m below the water's surface. Assume that the temperature of the air in the bubble remains constant as it rises. In terms of V_1 , calculate the volume of the bubble when it is just below the surface of the water.
 - If the air in the bubble cooled as it rose, the volume of the bubble would be less than the value calculated in part (c)(ii). Use physics principles to briefly explain why.

STOP

END OF EXAM