# AP<sup>®</sup> CALCULUS BC 2006 SCORING GUIDELINES

### Question 4

t (seconds)	0	10	20	30	40	50	60	70	80
v(t) (feet per second)	5	14	22	29	35	40	44	47	49

Rocket *A* has positive velocity v(t) after being launched upward from an initial height of 0 feet at time t = 0 seconds. The velocity of the rocket is recorded for selected values of *t* over the interval  $0 \le t \le 80$  seconds, as shown in the table above.

- (a) Find the average acceleration of rocket A over the time interval  $0 \le t \le 80$  seconds. Indicate units of measure.
- (b) Using correct units, explain the meaning of  $\int_{10}^{70} v(t) dt$  in terms of the rocket's flight. Use a midpoint

Riemann sum with 3 subintervals of equal length to approximate  $\int_{10}^{70} v(t) dt$ .

(c) Rocket *B* is launched upward with an acceleration of  $a(t) = \frac{3}{\sqrt{t+1}}$  feet per second per second. At time t = 0 seconds, the initial height of the rocket is 0 feet, and the initial velocity is 2 feet per second. Which of the two rockets is traveling faster at time t = 80 seconds? Explain your answer.

(a)	Average acceleration of rocket <i>A</i> is	1 : a	nswer
	$\frac{v(80) - v(0)}{80 - 0} = \frac{49 - 5}{80} = \frac{11}{20} \text{ ft/sec}^2$		
(b)	Since the velocity is positive, $\int_{10}^{70} v(t) dt$ represents the distance, in feet, traveled by rocket <i>A</i> from $t = 10$ seconds to $t = 70$ seconds.	3: {	<ul> <li>1 : explanation</li> <li>1 : uses v(20), v(40), v(60)</li> <li>1 : value</li> </ul>
	A midpoint Riemann sum is 20[v(20) + v(40) + v(60)] = 20[22 + 35 + 44] = 2020  ft		
(c)	Let $v_B(t)$ be the velocity of rocket <i>B</i> at time <i>t</i> . $v_B(t) = \int \frac{3}{\sqrt{t+1}} dt = 6\sqrt{t+1} + C$ $2 = v_B(0) = 6 + C$ $v_B(t) = 6\sqrt{t+1} - 4$ $v_B(80) = 50 > 49 = v(80)$	4 : {	$1: 6\sqrt{t+1}$ $1: constant of integration$ $1: uses initial condition$ $1: finds v_B(80)$ , compares to $v(80)$ ,and draws a conclusion
	Rocket <i>B</i> is traveling faster at time $t = 80$ seconds.		
Uni	ts of $ft/sec^2$ in (a) and ft in (b)	1 : u	nits in (a) and (b)

© 2006 The College Board. All rights reserved.

Visit apcentral.collegeboard.com (for AP professionals) and www.collegeboard.com/apstudents (for AP students and parents).









OR ALLOWED

CALCULUS AB SECTION II, Part B

Time-45 minutes

Number of problems—3

No calculator is allowed for these problems.

t (seconds)	0	10	20	30	40	50	60	70	80
v(t) (feet per second)	5	14	22	29	35	40	44	47	49



Continue problem 4 on page 11.

-10-



Δ

NO CALCULATOR ALLOWED

4

Work for problem 4(c)

$$\frac{R \circ c/kt}{a(t)} = \frac{3}{\sqrt{t+1}}$$

$$V(t) = \int_{V(t)} \frac{3}{\sqrt{t+1}} dt$$

$$\int_{U} \frac{1}{\sqrt{t+1}} dt$$

$$\int_{U} \frac{1}{\sqrt{t}} \frac{1}{\sqrt{t+1}} dt$$

$$\int_{U} \frac{1}{\sqrt{t}} \frac{1}{\sqrt{$$

Do not write beyond this border.

Rocket A (80) = 49 ft/sec

Rocket B is traveling faster at t=80 sec. Rocket B's velocity was found by v(t) = fa(t) dt and is 50 ft lsec. Rocket A's velocity was 49 ft/sec

### GO ON TO THE NEXT PAGE.

© 2006 The College Board. All rights reserved. Visit apcentral.collegeboard.com (for AP professionals) and www.collegeboard.com/apstudents (for students and parents).







## - CALCULUS AB

#### SECTION II, Part B

Time—45 minutes

Number of problems—3

No calculator is allowed for these problems.

·			$\frown$	$\wedge$		$ \land $	$\sim$	$\sim$	
t (seconds)	0	10	20	30	40	50	60	70	80
v(t) (feet per second)	5	14	22	29	35	. 40	44	47	49
	V	$\overline{\langle}$		$\overline{\checkmark}$					

Work for problem 4(a) $\frac{1}{80}\left(9+8+7+6+5+4+3+2\right)=$  $=\frac{46}{80}=\frac{23}{40}$  H/sec<sup>2</sup> Do not write beyond this border. WING DEADIN THIS DOINE Work for problem 4(b) S v(t) at indicates the distance the rocket traveled over 10 the interval 10st = 70. In this case, it indicates total distance because the rocket's velocity is only increasing during this time period. (22+2(35)+44)=136 ft.

Continue problem 4 on page 11.

B

-10-

© 2006 The College Board. All rights reserved.

Visit apcentral.collegeboard.com (for AP professionals) and www.collegeboard.com/apstudents (for students and parents).

Work for problem 4(c)3(+H)<sup>1/2</sup> V(+)=61++1+C 2=6 0+1+0  $7 = l_0 + c$ -4=0 v(t) = 6 + 1 - 4Do not write beyond this border V(8)=6 J80H -4 6.9-4 V(80)=52 ft/sec Rocket B is traveling faster at += 80. The table states that Pocket A is traveling at 49 ft/sec. By taking the antideravitive for alt) of Rocket B and solving with initial Conditions for C, then substituting += 80, We find V(80)=52 Ft/sec, faster than Rocket A.

NO CALCULATOR ALLOWED

GO ON TO THE NEXT PAGE.

4B,

© 2006 The College Board. All rights reserved. Visit apcentral.collegeboard.com (for AP professionals) and www.collegeboard.com/apstudents (for students and parents).







CALCULUS AB

**SECTION II, Part B** 

Time—45 minutes

Number of problems----3

No calculator is allowed for these problems.

t (seconds)	0	10	20	30	40	50	60	70	80
v(t) (feet per second)	5	14	22	29	35	40	44	47	49

Work for problem 4(a)acceleration are = 11 feet / second Do not write beyond this border. ביט ווטר איוויב טכץטווע נונוא טטועכן. The integral of the velocity is the position. Sio v(t) dt means the position of the rocket from 10 sec to 70 sec. Work for problem 4(b)

Continue problem 4 on page 11.

-10-

© 2006 The College Board. All rights reserved.

Visit apcentral.collegeboard.com (for AP professionals) and www.collegeboard.com/apstudents (for students and parents).



GO ON TO THE NEXT PAGE.

© 2006 The College Board. All rights reserved. Visit apcentral.collegeboard.com (for AP professionals) and www.collegeboard.com/apstudents (for students and parents).

Do not write beyond this border.

# AP<sup>®</sup> CALCULUS BC 2006 SCORING COMMENTARY

## **Question 4**

## Overview

This problem presented students with a table of velocity values for rocket *A* at selected times. In part (a) students needed to recognize the connection between the average acceleration of the rocket over the given time interval and the average rate of change of the velocity over this interval. In part (b) students had to recognize the definite integral as the total change, in feet, in rocket *A*'s position from time t = 10 seconds to time t = 70 seconds and then approximate the value of this definite integral using a midpoint Riemann sum and the data in the table. Units of measure were important in both parts (a) and (b). Part (c) introduced a second rocket and gave its acceleration in symbolic form. The students were asked to compare the velocities of the two rockets at time t = 80 seconds. The velocity of rocket *B* could be determined by finding the antiderivative of the acceleration and using the initial condition or by using the Fundamental Theorem of Calculus and computing a definite integral.

Sample: 4A Score: 9

The student earned all 9 points.

## Sample: 4B Score: 6

The student earned 6 points: 2 points in part (b), 3 points in part (c), and the units point. The first line in part (a) is a correct but uncommon method. An error was made in the addition. In part (b) the explanation is acceptable; the student correctly identifies the midpoint values but the method is incorrect. In part (c) the fourth point was not earned due to an error in the computation of v(80) for rocket *B*.

## Sample: 4C Score: 3

The student earned 3 points: 1 point in part (a) and 2 points in part (c). The units are correct in part (a) but missing in part (b). For the explanation in part (b), the integral does not represent the position but rather the displacement over the time interval [10, 70]. The constant of integration does not appear in part (c) so the student earned the antiderivative and comparison points only.