
AP® CALCULUS BC 2010 SCORING GUIDELINES (Form B)

Question 1

In the figure above, R is the shaded region in the first quadrant bounded by the graph of $y = 4\ln(3-x)$, the horizontal line y = 6, and the vertical line x = 2.



- (a) Find the area of R.
- (b) Find the volume of the solid generated when R is revolved about the horizontal line y = 8.
- (c) The region R is the base of a solid. For this solid, each cross section perpendicular to the x-axis is a square. Find the volume of the solid.
 - 1 : Correct limits in an integral in (a), (b), or (c)

(a)
$$\int_0^2 (6 - 4\ln(3 - x)) dx = 6.816$$
 or 6.817

 $2:\begin{cases} 1 : integrand \\ 1 : answer \end{cases}$

(b)
$$\pi \int_0^2 ((8 - 4\ln(3 - x))^2 - (8 - 6)^2) dx$$

= 168.179 or 168.180

 $3: \begin{cases} 2: integrand \\ 1: answer \end{cases}$

(c)
$$\int_0^2 (6 - 4\ln(3 - x))^2 dx = 26.266$$
 or 26.267


 $3: \begin{cases} 2: \text{integrand} \\ 1: \text{answer} \end{cases}$

CALCULUS BC SECTION II, Part A

Time-45 minutes

Number of problems—3

A graphing calculator is required for some problems or parts of problems.



梦

Work for problem 1(a)

$$SR = 2 \times 6 - \int_{0}^{2} 4 \ln (3 \times) dx$$

= $2 \times 12 - (5.183) = 6.817$

Work for problem 1(b)

assum
$$y_1 = 4 \ln (3-x)$$

 $y_2 = 6$
 $V = \pi \sqrt{\frac{(8-y_1)^2}{(8-y_1)^2}} \frac{(8-y_2)^2}{4x}$
 $= \pi \sqrt{\frac{(8-y_1)^2}{(8-4\ln(3+x))^2-(8-6)^2}} dx$
 $\approx (68.180)$

Work for problem 1(c)

Do not write beyond this border.

$$V = \int_{0}^{2} \left[6 - 4 \ln (3-x)\right]^{2} dx$$

 ≈ 26.267


דים ווחר אוזוכ הבלחוום חווי החוחבוי

CALCULUS AB SECTION II, Part A

Time—45 minutes

Number of problems -3

A graphing calculator is required for some problems or parts of problems.



Work for problem 1(a)

Do not write beyond this border.

$$R = \int_{0}^{2} 6 - 4 \ln(3 - x) dx$$

$$= \int_{0}^{2} 6 dx - \int_{0}^{2} 4 \ln(3 - x) dx$$

$$= 6x/_{0}^{2} - 4 \int_{0}^{2} \ln(3 - x) dx$$

$$f = \ln(3 - x) \quad f' = \frac{1}{3 - x}$$

$$g' = 1 \quad g = x$$

$$= 12 - \left(4x \ln(3 - x)/_{0}^{2} - \int_{0}^{2} \frac{-x}{3 - x} dx\right)$$

$$= 6.817 \text{ units}^{2}$$

Continue problem 1 on page 5.

1 1 1 1 1 1 1 1 1 1

Work for problem 1(b)

$$V = T \int_{0}^{2} [8-4\ln(3-x)]^{2} - (8-6)^{2} dx$$

$$= T \int_{0}^{2} [8-4\ln(3-x)]^{2} - 2^{2} dx$$

$$= 168. [80 units^{3}]$$

Work for problem 1(c)

$$V = \int_{0}^{2} (2x6)^{2} - (2x4\ln(3-x)^{2}) dy$$

$$= \int_{0}^{2} (2^{2} - (8\ln(3-x))^{2}) dx$$

$$= 222. [33] \text{ units}^{3}$$


GO ON TO THE NEXT PAGE.

CALCULUS BC SECTION II, Part A

Time-45 minutes

Number of problems—3

A graphing calculator is required for some problems or parts of problems.



Work for problem 1(a)

 $A = \int_0^2 (6-4\ln(3-x)) dx$ with the help of the calculator, we type this in and we get $A = \int_0^2 (6-4\ln(3-x)) dx = 6.81$ Work for problem 1(b)

Since the p is revolved about l: y=1 we use the washer method $V=\pi \int_0^2 (8-6^2)(8-4\ln(3-x)^2) dx$ With the calculator, we get $V=\pi \int_0^2 (8-6^2)(8-(\alpha\ln(3-x))^2) dx = (1.05)^2$

Work for problem 1(c)

Do not write beyond this border.

The side of the square shall be $y = 4 \ln (3-x)$ which makes the Area of the square as $A = (4 \ln (3-x))^2$ $V = \pi \int_0^x (4 \ln (3-x))^2 dx$ We use the calculator to get $V = \pi \int_0^x (4 \ln (3-x))^2 dx = 51.73x$

AP® CALCULUS BC 2010 SCORING COMMENTARY (Form B)

Question 1

Sample: 1A Score: 9

The student earned all 9 points.

Sample: 1B Score: 6

The student earned 6 points: the global limits point, 2 points in part (a), 3 points in part (b), and no points in part (c). In part (a) the student earned both points and the global limits point. The student's intermediate work includes a misplaced 4, but the correct numerical answer was treated as a restart since this was the calculator portion of the exam. In part (b) the student's work is correct. In part (c) the student does not use square cross sections and was not eligible for any points.

Sample: 1C Score: 3

The student earned 3 points: the global limits point, 2 points in part (a), no points in part (b), and no points in part (c). In part (a) the student earned the global limits point and has correct work. In part (b) the student attempts to find the volume using washers, but the work is incorrect. In part (c) the student uses an incorrect width for the area of the square cross section and includes a factor of π .