

AP Physics C: Mechanics 2000 Student Samples

The materials included in these files are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 3,900 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT®, the PSAT/NMSQT™, the Advanced Placement Program® (AP®), and Pacesetter®. The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.


мммммммммм⁶¹

Mech 2.

A rubber ball of mass m is dropped from a cliff. As the ball falls, it is subject to air drag (a resistive force caused by the air). The drag force on the ball has magnitude bv^2 , where b is a constant drag coefficient and v is the instantaneous speed of the ball. The drag coefficient b is directly proportional to the cross-sectional area of the ball and the density of the air and does not depend on the mass of the ball. As the ball falls, its speed approaches a constant value called the terminal speed.

(a) On the figure below, draw and label all the forces on the ball at some instant before it reaches terminal speed.

Fig. = Juag Force




(b) State whether the magnitude of the acceleration of the ball of mass *m* increases, decreases, or remains the same as the ball approaches terminal speed. Explain.

The magnitude of the acceleration decreases as the hall approaches terminal speed because the drag force will increase until it is equal to the force due to gravity. Once Fig. = mg. The ball will be at terminal speed because the acceleration will be zero.

(c) Write, but do NOT solve, a differential equation for the instantaneous speed v of the ball in terms of time r, the given quantities, and fundamental constants.

(d) Determine the terminal speed v_t in terms of the given quantities and fundamental constants.



(c) Determine the energy dissipated by the drag force during the fall if the ball is released at height h and teaches its terminal speed before hitting the ground, in terms of the given quantities and fundamental constants.



Energy dissipated =
$$E_A - E_B$$

= $mgh - \frac{1}{2}mv_E^2$
= $mgh - \frac{1}{2}m(\frac{mg}{6})$
= $mg(h - \frac{1}{2}\frac{m}{6})$

M M M M M M M M M M M M #1

Mech 2,

A subber ball of mass m is dropped from a cliff. As the ball falls, it is subject to air drag (a resistive force caused by the air). The drag force on the ball has magnitude bv^2 , where b is a constant drag coefficient and v is the instantaneous speed of the ball. The drag coefficient b is directly proportional to the cross-sectional area of the ball and the density of the air and does not depend on the mass of the ball. As the ball falls, its speed approaches a constant value called the terminal speed.

(a) On the figure below, draw and label all the forces on the ball at some instant before it reaches terminal speed.

1 mg

(b) State whether the magnitude of the acceleration of the ball of mass *m* increases, decreases, or remains the same as the ball approaches terminal speed. Explain, $\sqrt{g - b \sqrt{2}} = a - g - \sqrt{\frac{\sqrt{m} \frac{3}{2}}{b}} = a$

increases

Let's compare $g = bx^2$ for a \$ g - gy = aIn the first equation assigner $g - by^2 = a$ number is being suptracted from $g - by^2 = a$

(c) Write, but do NOT solve, a differential equation for the instantaneous speed v of the ball in terms of time t, the given quantities, and fundamental constants.

mg=ma+bv2
mg=mdy+bv2
mg=mdy+bv2
mg=mdy+bv2

mg=mdy+bv2

M M M M M M M M M M M M

#2

(d) Determine the terminal speed v_t in terms of the given quantities and fundamental constants.

$$mg - bv^2 = moe$$

$$mg = bv^2$$

$$mg = v^2$$

$$v = \sqrt{mg}$$


(e) Determine the energy dissipated by the drag force during the fall if the ball is released at height h and reaches its terminal speed before hitting the ground, in terms of the given quantities and fundamental constants.

M M M M M M M M M M M M 11

Mech 2.

A rubber ball of mass m is dropped from a cliff. As the ball falls, it is subject to air drag (a resistive force caused by the air). The drag force on the ball has magnitude bv^2 , where b is a constant drag coefficient and v is the instantaneous speed of the ball. The drag coefficient b is directly proportional to the cross-sectional area of the ball and the density of the air and does not depend on the mass of the ball. As the ball falls, its speed approaches a constant value called the terminal speed.

(a) On the figure below, draw and label all the forces on the ball at some instant before it reaches terminal speed.



(b) State whether the magnitude of the acceleration of the ball of mass m increases, decreases, or remains the same as the ball approaches terminal speed. Explain.

The acceleration decreases we to the drag force.

(c) Write, but do NOT solve, a differential equation for the instantaneous speed v of the ball in terms of time t, the given quantities, and fundamental constants.

$$mg - bv^{2} = ma$$


$$mg - bv^{2} = ma$$

$$\left(\frac{ma}{m} - \frac{bv^{2}}{m}\right)dt = dV$$

$$\int \left(g - \frac{bv^{2}}{m}\right) = V$$

M M M M M M M M M M M M 12

(d) Determine-the terminal speed v_i in terms of the given quantities and fundamental constants.



$$mg - bv^2 = mq$$
 $mg - bv^2 = mqv$
 dt

(e) Determine the energy dissipated by the drag force during the fall if the ball is released at height h and reaches its terminal speed before hitting the ground, in terms of the given quantities and fundamental constants.