

AP Statistics 2000 Student Samples

The materials included in these files are intended for non-commercial use by AP teachers for course and exam preparation; permission for any other use must be sought from the Advanced Placement Program. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.

These materials were produced by Educational Testing Service (ETS), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 3,900 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT®, the PSAT/NMSQT™, the Advanced Placement Program® (AP®), and Pacesetter®. The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

(a) Based on the scatterplot, describe the effect of drug A and how it is related to strength in milligrams.

Drug A gives a post surgical pain relief of about 60 to 70 (on a reale 0-100). The amount of relief is not affected by the strength in milligrams

(b) Based on the scatterplot, describe the effect of drug B and how it is related to strength in milligrams.

Drug B has little effect on pain relief for about 210-270 mg dosages. The amount of relief then increases linearly with strength (in mg) reaching a maximum relief of about 90 for a 400 mg dosage

(c) Which drug would you give and at what strength, if the goal is to get pain relief of at least 50 at the lowest possible strength? Justify your answer based on the scatterplot.

I would give Drug A at a strength of 210 mg.

Drug A is consistently above 50 in relief and it
is essentially independent of strength, so one can
use small dosages. The smallest dosage of Drug

B that one can use for a relief of above 50,
on the other hand, is about 330 mg

(a) Based on the scatterplot, describe the effect of drug A and how it is related to strength in milligrams.

Drug A seems to give the postsurgical patients à pain relief rating of 50 to 70, regardless of the dosage amount. Pain relief with drug A and the dosage strength have a linear relationship with a slope of approx. O, meaning changing the strength of the dosage does not change pain relief.

(b) Based on the scatterplot, describe the effect of drug B and how it is related to strength in milligrams.

Drug B seems to bring more pain relief with higher dores. Pain relief with Drug B and the dorage strength have a positive linear relationship which means they increase with each other. The higher the dosage of Drug B, the more pain relief there is:

(c) Which drug would you give and at what strength, if the goal is to get pain relief of at least 50 at the lowest possible strength? Justify your answer based on the scatterplot.

By looking at the scatterplot, one can see that:

- Drug A consistently gives a pair relief level of 60, from 210 to 400 mg.

- Drug. B murt be given at a 325 mg. dorage to provide level 50 velief.

Therefore, to give the least medication and achieve a relief level of at least 59 give drug A at the lowert dwage, 210 mg, to give relief of at least 50 w lear mg of medication.

(a) Based on the scatterplot, describe the effect of drug A and how it is related to strength in milligrams.

Because the relationship is almost horizontal, the effect of drug A remains about the same with increasing strength (from 200 mg to 400 mg). Drug A is known to relieve about 65% of the pain at any strength.

(b) Based on the scatterplot, describe the effect of drug B and how it is related to strength in milligrams.

Drug B's effectiveness in pain relief and its strength have a positive relationship. It is almost ineffective from about 270 mg but its effectiveness in pain relief increases as strength increases thereafter. At 400 mg, the pain relief is about 90%.

(c) Which drug would you give and at what strength, if the goal is to get pain relief of at least 50 at the lowest possible strength? Justify your answer based on the scatterplot.

I would give drug A at 210 milligrams given the preconditions. Drug A has the lowest strength at which at least 50% of the pain is relieved (2004 65%) and thus is the best choice.